

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Anti-Tyrosinase-Related Protein-1 (TYRP-1) (Melanoma Marker) Monoclonal Antibody(Clon: TYRP1/1986), Clon: [TYRP1/1986], Mouse ABI-36-3335

|                            |  |
|----------------------------|--|
| Article Name               | Anti-Tyrosinase-Related Protein-1 (TYRP-1) (Melanoma Marker)<br>Monoclonal Antibody(Clon: TYRP1/1986), Clon: [TYRP1/1986],<br>Mouse  |
| Biozol Catalog Number      | ABI-36-3335  |
| Supplier Catalog Number    | 36-3335  |
| Alternative Catalog Number | ABI-36-3335-100UG  |
| Manufacturer               | Abeomics   |
| Host                       | Mouse  |
| Category                   | Antikörper   |
| Application                | ELISA, IHC   |
| Species Reactivity         | Human  |
| Immunogen                  | A recombinant fragment (around aa 257-377) of human TYRP1<br>protein (exact sequence is proprietary)   |
| Product Description        | It reacts with a 75kDa melanocyte-specific gene product, identified<br>as Tyrosinase-related protein-1 (TRP-1). It is involved in melanin<br>synthesis. TRP1 is present on the melanosomal membranes of<br>melanoma, normal melanocytes and nevi.Recent evidence su... |
| Clonality                  | Monoclonal   |
| Clone Designation          | [TYRP1/1986]   |
| NCBI                       | <a href="#">7306</a>   |

|                    |  |
|--------------------|--|
| UniProt            | <a href="#">P17643</a>   |
| Application Dilute | ELISA (Use Ab at 2-4ug/ml for coating) (Order Ab without BSA),<br>,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 |