

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

PECAM-1 Antibody / CD31, Clone: [C31.7], Unconjugated, Mouse, Monoclonal BYT-ORB2637883

Article Name	PECAM-1 Antibody / CD31, Clone: [C31.7], Unconjugated, Mouse, Monoclonal
Biozol Catalog Number	BYT-ORB2637883
Supplier Catalog Number	orb2637883
Alternative Catalog Number	BYT-ORB2637883-100
Manufacturer	Biorbyt
Host	Mouse
Category	Antikörper
Application	FACS, IHC-P, WB
Species Reactivity	Human
Immunogen	Human recombinant PECAM-1 / CD31 protein was used as the immunogen for this antibody.
Conjugation	Unconjugated
Product Description	PECAM-1, or platelet endothelial cell adhesion molecule-1, also called CD31, is made up of six extracellular immunoglobulin-like domains, a single transmembrane domain, and a cytoplasmic domain containing serine and tyrosine residues that when phosph...
Clonality	Monoclonal
Clone Designation	[C31.7]
Buffer	1 mg/ml in 1X PBS, rAlbumin free, sodium azide free

Application Dilute	Flow cytometry: 1-2ug/10 ⁶ cells,Western blot: 1-2ug/ml,Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Application Notes	<p>Application Notes: The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the PECAM-1 antibody to be titered up or down for optimal performance.</p> <ol style="list-style-type: none">1. Staining of formalin-fixed tissues requires boiling tissue sections in 1mM EDTA, pH 7.5-8.5, for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min