

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Cytokeratin 18 Antibody, Clone: [KRT18/835], Unconjugated, Mouse, Monoclonal BYT-ORB2638774**

Article Name	Cytokeratin 18 Antibody, Clone: [KRT18/835], Unconjugated, Mouse, Monoclonal
Biozol Catalog Number	BYT-ORB2638774
Supplier Catalog Number	orb2638774
Alternative Catalog Number	BYT-ORB2638774-100
Manufacturer	Biorbyt
Host	Mouse
Category	Antikörper
Application	IHC-P, WB
Species Reactivity	Human, Rat
Immunogen	Recombinant full-length human KRT18 protein was used as the immunogen for the Cytokeratin 18 antibody.
Conjugation	Unconjugated
Product Description	This mAb reacts with a wide variety of simple epithelia. It does not react with stratified squamous epithelia. It reacts with epithelial tumors of the gastrointestinal tract, lung, breast, pancreas, ovary, and thyroid. Cytokeratin 18, which belongs t...
Clonality	Monoclonal
Clone Designation	[KRT18/835]
UniProt	<a href="#">P05783</a>

Buffer	1 mg/ml in 1X PBS, rAlbumin free, sodium azide free
Application Dilute	Western blot: 1-2ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Application Notes	Application Notes: Optimal dilution of the Cytokeratin 18 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min