BIOZOL

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit IgG anti-Sheep IgG (F(ab)2)-unconj., MinX none

DNA-SEC-182618

Article Name	Rabbit IgG anti-Sheep IgG (F(ab)2)-unconj., MinX none
Biozol Catalog Number	DNA-SEC-182618
Supplier Catalog Number	SEC-182618
Alternative Catalog Number	DNA-SEC-182618
Manufacturer	dianova
Host	Rabbit
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Sheep
Immunogen	Sheep IgG F(ab)2 fragment
Conjugation	Unconjugated
Format	IgG
Target Specificity	IgG (F(ab')2)
Cross-Adsorption (MinX)	no cross-adsorbtion

Product Description	Anti-Sheep IgG F(ab)2 Antibody generated in rabbit recognizes the dimeric Fab portion of the sheep IgG molecule. Sheep IgG F(ab)2 is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme pepsin under controlle...
Clonality	Polyclonal
Concentration	10.0 mg/mL
Isotype	Ig
Buffer	0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Sheep IgG, Sheep IgG F(ab')2 and Sheep Serum. No reaction was observed against Sheep IgG F(c).
Form	Lyophilized
Formula	10 mM NaPO4,150 mM NaCl,pH 7,2,lyophilisate,Azide/BSA free
Target	Sheep
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000
Application Notes	Anti-Sheep IgG F(ab)2 antibody is suitable for use in ELISA, immunohistochemistry, and western blot. Specific conditions for reactivity should be optimized by the end user.