BIOZOL

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit F(ab)2 anti-Sheep IgG (F(ab)2)-HRPO, MinX none

DNA-SEC-182718

Article Name	Rabbit F(ab)2 anti-Sheep IgG (F(ab)2)-HRPO, MinX none
Biozol Catalog Number	DNA-SEC-182718
Supplier Catalog Number	SEC-182718
Alternative Catalog Number	DNA-SEC-182718
Manufacturer	dianova
Host	Rabbit
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Sheep
Immunogen	Sheep IgG F(ab)2 fragment
Conjugation	HRPO
Format	F(ab')2
Target Specificity	IgG (F(ab')2)
Cross-Adsorption (MinX)	no cross-adsorbtion

Product Description	F(ab)2 Anti-Sheep IgG Fab2 Peroxidase Antibody generated in rabbit detects immunoglobulin g from sheep. Each IgG has two antigen binding sites. Representing approximately 75% of serum immunoglobulins in humans, IgG is the most abundant antibody isoty...
Clonality	Polyclonal
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product is a F(ab')2 fragment of an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation, ion exchange chromatography and pepsin digestion followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum, Sheep IgG, Sheep IgG F(ab')2 and Sheep Serum. No reaction was observed against Sheep IgG F(c), anti-Rabbit IgG F(c) or anti-Pepsin.
Form	Lyophilized
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin
Target	Sheep
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:10,000 - 1:50,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, Western Blot Dilution: 1:1,000 - 1:10,000
Application Notes	F(ab)2 Anti-Sheep IgG Fab2 Peroxidase secondary antibody reagents are ideal for ELISA, western blotting, Immunohistochemistry, Fluorescence Microscopy, Flow Cytometry as well as other antibody detection methods.