**BIOZOL**

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

**E-Mail:** info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

# Product Datasheet

## Mouse IgG anti-Goat IgG (H+L)-unconj., MinX none

## DNA-SEC-182835

| | |
|---|---|
| Article Name | Mouse IgG anti-Goat IgG (H+L)-unconj., MinX none |
| Biozol Catalog Number | DNA-SEC-182835 |
| Supplier Catalog Number | SEC-182835 |
| Alternative Catalog Number | DNA-SEC-182835 |
| Manufacturer | dianova |
| Host | Mouse |
| Category | Antikörper |
| Application | ELISA,IHC,WB |
| Species Reactivity | Goat |
| Immunogen | Goat IgG whole molecule |
| Conjugation | Unconjugated |
| Format | IgG |
| Target Specificity | IgG (H+L) |
| Cross-Adsorption (MinX) | no cross-adsorbtion |

| | |
|---|---|
| Product Description | Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level o... |
| Clonality | Polyclonal |
| Concentration | 1.0 mg/ml |
| Isotype | Ig |
| Buffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Purity | Goat IgG (H&L) Antibody was prepared from monospecific polyclonal ascites by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Mouse Serum, Goat IgG and Goat Serum. No reaction was observed against Rabbit, Human or Mouse Serum Proteins. |
| Form | Liquid (sterile filtered) |
| Formula | 20 mM K3PO4,150 mM NaCl,pH 7,2,sterile filtered,0,01% NaN3 |
| Target | Goat |
| Antibody Type | Secondary Antibody |
| Application Dilute | ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000 |
| Application Notes | Anti-Goat IgG antibody has been tested by ELISA and is suitable for ELISA, western blot, and immunohistochemistry, as well as other assays requiring lot-to-lot consistency. |