BIOZOL

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit IgG anti-Sheep IgG (H+L)-HRPO, MinX none

DNA-SEC-183548

Article Name	Rabbit IgG anti-Sheep IgG (H+L)-HRPO, MinX none
Biozol Catalog Number	DNA-SEC-183548
Supplier Catalog Number	SEC-183548
Alternative Catalog Number	DNA-SEC-183548
Manufacturer	dianova
Host	Rabbit
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Sheep
Immunogen	Anti-Sheep IgG (H&L) was produced by repeated immunization with Sheep IgG whole molecule in rabbit.
Conjugation	HRPO
Format	IgG
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	no cross-adsorbtion

Product Description	Anti-Sheep IgG antibody generated in rabbit detects specifically sheep IgG. This secondary antibody anti-Sheep is ideal for investigators who routinely perform ELISA, Sandwich ELISA, titration assays, western-blot, immunoprecipitation and more genera...
Clonality	Polyclonal
Concentration	2.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Sheep IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum, Sheep IgG and Sheep Serum.
Form	Lyophilized
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin
Target	Sheep
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:40,000 - 1:80,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, Western Blot Dilution: 1:1,000 - 1:5,000
Application Notes	Anti-Sheep IgG (HRP) is suitable for use in immunoelectrophoresis, western-blot, competitive western-blot, ELISA and competitive ELISA assays. Specific conditions for reactivity and signal detection should be optimized by the end user.