

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Sheep anti Human IgD (Fc specific), Clone: [Polyclonal], Monoclonal NMB-SAHU/IGD(FC)

|                            |   |
|----------------------------|---|
| Article Name               | Sheep anti Human IgD (Fc specific), Clone: [Polyclonal], Monoclonal   |
| Biozol Catalog Number      | NMB-SAHU/IGD(FC)  |
| Supplier Catalog Number    | ShAHu/IgD(Fc)   |
| Alternative Catalog Number | NMB-SAHU/IGD(FC)  |
| Manufacturer               | NordicMubio   |
| Host                       | Sheep   |
| Category                   | Antikörper  |
| Species Reactivity         | Human   |
| Conjugation                | Unconjugated  |
| Format                     | Antiserum   |
| Target Specificity         | IgD   |
| Cross-Adsorption (MinX)    | no cross-adsorbtion   |
| Product Description        | The reactivity of the antiserum is restricted to the Fc part of the IgD molecule. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against appropriate concentrations of the immunogen, a single precipitin lin... |
| Clonality                  | Monoclonal  |
| Clone Designation          | [Polyclonal]  |

|                   |  |
|-------------------|--|
| Buffer            | Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal sheep serum. No foreign proteins added. Reconstitute the lyophilised   |
| Source            | Highly purified homogenous IgD isolated from human serum. Freunds complete adjuvant is used in the first step of the immunization procedure.   |
| Formula           | Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal sheep serum. No foreign proteins added.  |
| Antibody Type     | Secondary Antibody   |
| Application Notes | Precipitation assays. The lyophilized antiserum is shipped at ambient temperature and may be stored at +4C, prolonged storage at or below -20C. In immunoelectrophoresis use 2 µl serum or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in 3 mm diameter center well and 2 µl serum samples (neat and serially diluted) in 2 mm diameter peripheral wells. |