

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Sheep Antiserum anti-Rat IgG2b (Fc)-unconj., MinX none NMB-SHARA/IGG2B

Article Name	Sheep Antiserum anti-Rat IgG2b (Fc)-unconj., MinX none
Biozol Catalog Number	NMB-SHARA/IGG2B
Supplier Catalog Number	ShARa/IgG2b
Alternative Catalog Number	NMB-SHARA/IGG2B
Manufacturer	NordicMubio
Host	Sheep
Category	Antikörper
Species Reactivity	Rat
Conjugation	Unconjugated
Format	Antiserum
Target Specificity	IgG2b (Fc)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	The reactivity of the antiserum is directed to the Fc portion of the subclass IgG2b. It does not react with other subclasses of IgG, IgG/Fab fragments, IgM and IgA or any non-Ig protein in rat serum, as tested by immunoelectrophoresis and double radi
Clonality	Polyclonal
Clone Designation	[Polyclonal]

Buffer	Delipidated, heat inactivated, lyophilized whole antiserum. No preservative added as it may interfere with he antibody activity. Total protein and IgG concentration in the antiserum are comparable to those of pooled normal sheep serum. No foreign protein
Source	Pools of purified homogenous IgG2b isolated from pooled rat serum. Freunds complete adjuvant is used in the first step of the immunization procedure.
Formula	Delipidated, heat inactivated, lyophilized whole antiserum. No preservative added as it may interfere with he antibody activity. Total protein and IgG concentration in the antiserum are comparable to those of pooled normal sheep serum. No foreign protein
Antibody Type	Secondary Antibody
Application Notes	Precipitation assays. In immunoelectrophoresis use 2 μ l or equivalent against 120 μ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 μ l antiserum in a 3 mm diameter centre well and 2 μ l serum samples (neat and