

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Swine Antiserum anti-Mouse IgA-unconj., MinX none NMB-SWAM/IGA(FC)

Article Name	Swine Antiserum anti-Mouse IgA-unconj., MinX none
Biozol Catalog Number	NMB-SWAM/IGA(FC)
Supplier Catalog Number	SwAM/IgA(Fc)
Alternative Catalog Number	NMB-SWAM/IGA(FC)
Manufacturer	NordicMubio
Host	Porcine
Category	Antikörper
Species Reactivity	Mouse
Conjugation	Unconjugated
Format	Antiserum
Target Specificity	IgA
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	The reactivity of the antiserum is restricted to the Fc part of the IgA molecule. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against normal mouse serum a single precipitin line is obtained which shows
Clonality	Polyclonal
Clone Designation	[Polyclonal]

Buffer	Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal swine serum. No foreign proteins added. Reconstitute the lyophilized
Source	Highly purified pools of monoclonal IgA isolated from mouse serum. Freunds complete adjuvant is used in the first step of the immunization procedure.
Formula	Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal swine serum. No foreign proteins added.
Antibody Type	Secondary Antibody
Application Notes	Precipitation assays. In immunoelectrophoresis use 2 $\mu$ l serum or equivalent against 120 $\mu$ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 $\mu$ l antiserum in 3 mm diameter center well and 2 $\mu$ l serum samples (neat