

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

**Swine anti Rabbit Ig light chains kappa and lambda (free and bound),
Clone: [Polyclonal], Porcine, Monoclonal
NMB-SWAR/L(SD+HD)**

Article Name	Swine anti Rabbit Ig light chains kappa and lambda (free and bound), Clone: [Polyclonal], Porcine, Monoclonal
Biozol Catalog Number	NMB-SWAR/L(SD+HD)
Supplier Catalog Number	SwAR/L(SD+HD)
Alternative Catalog Number	NMB-SWAR/L(SD+HD)
Manufacturer	NordicMubio
Host	Porcine
Category	Antikörper
Product Description	As tested in immunoelectrophoresis and double radial immunodiffusion against pooled normal rabbit serum and purified IgG fractions and light chain preparations, the antiserum reacts with determinants exposed on the surface (SD) and the Fab portion of...
Clonality	Monoclonal
Clone Designation	[Polyclonal]
Buffer	Delipidated, heat inactivated lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled swine ser
Source	Purified light chains isolated from polyclonal rabbit IgG. Freunds complete adjuvant is used in the first step of the immunization procedure.

Formula	<p>Delipidated, heat inactivated lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled swine se</p>
Application Notes	<p>Precipitation assays. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and serially diluted) in 2 mm diameter peripheral wells. In single immunodiffusion, use 1% antiserum in agar gel to establish a standard curve.</p>