

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Hu CD20 Purified Azide Free, Clone: [MEM-97], Monoclonal EXB-10-217-C025

Article Name Anti-Hu CD20 Purified Azide Free, Clone: [MEM-97], Monoclonal Biozol Catalog Number EXB-10-217-C025 Supplier Catalog Number 10-217-C025 Alternative Catalog Number EXB-10-217-C025 Manufacturer EXBIO Category Antikörper Application FC, IP Species Reactivity Porcine, Human, Bovine Immunogen Raji human Burkitts lymphoma cell line Product Description CD20 is a cell surface 33-37 (depending on the degree of phosphorylation) kDa non-glycosylated surface phosphorprotein expressed on mature and most malignant B cells, but not stem cells or plasma cells (low number of the CD20 has been also detected on Clonality Monoclonal Concentration 1 mg/ml Clone Designation [MEM-97] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4 Storage 2°C to 8°C		
Supplier Catalog Number 10-217-C025 Alternative Catalog Number EXB-10-217-C025 Manufacturer EXBIO Category Antikörper Application FC, IP Species Reactivity Porcine, Human, Bovine Immunogen Raji human Burkitts lymphoma cell line CD20 is a cell surface 33-37 (depending on the degree of phosphorylation) kDa non-glycosylated surface phosphoprotein expressed on mature and most malignant B cells, but not stem cells or plasma cells (low number of the CD20 has been also detected on Clonality Monoclonal Concentration 1 mg/ml Clone Designation [MEM-97] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4	Article Name	Anti-Hu CD20 Purified Azide Free, Clone: [MEM-97], Monoclonal
Alternative Catalog Number EXB-10-217-C025 Manufacturer EXBIO Category Antikörper Application FC, IP Species Reactivity Porcine, Human, Bovine Immunogen Raji human Burkitts lymphoma cell line CD20 is a cell surface 33-37 (depending on the degree of phosphorylation) kDa non-glycosylated surface phosphoprotein expressed on mature and most malignant B cells, but not stem cells or plasma cells (low number of the CD20 has been also detected on Clonality Monoclonal Concentration 1 mg/ml Clone Designation [MEM-97] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4	Biozol Catalog Number	EXB-10-217-C025
Manufacturer EXBIO Category Antikörper Application FC, IP Species Reactivity Porcine, Human, Bovine Immunogen Raji human Burkitts lymphoma cell line CD20 is a cell surface 33-37 (depending on the degree of phosphorylation) kDa non-glycosylated surface phosphoprotein expressed on mature and most malignant B cells, but not stem cells or plasma cells (low number of the CD20 has been also detected on Clonality Monoclonal Concentration 1 mg/ml Clone Designation [MEM-97] Isotype IgG1 Phosphate buffered saline (PBS), pH 7.4	Supplier Catalog Number	10-217-C025
Category Antikörper Application FC, IP Species Reactivity Porcine, Human, Bovine Immunogen Raji human Burkitts lymphoma cell line CD20 is a cell surface 33-37 (depending on the degree of phosphorylation) kDa non-glycosylated surface phosphoprotein expressed on mature and most malignant B cells, but not stem cells or plasma cells (low number of the CD20 has been also detected on Clonality Monoclonal Concentration 1 mg/ml Clone Designation [MEM-97] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4	Alternative Catalog Number	EXB-10-217-C025
Application FC, IP Species Reactivity Porcine, Human, Bovine Immunogen Raji human Burkitts lymphoma cell line CD20 is a cell surface 33-37 (depending on the degree of phosphorylation) kDa non-glycosylated surface phosphoprotein expressed on mature and most malignant B cells, but not stem cells or plasma cells (low number of the CD20 has been also detected on Clonality Monoclonal Concentration 1 mg/ml Clone Designation [MEM-97] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4	Manufacturer	EXBIO
Species Reactivity Porcine, Human, Bovine Raji human Burkitts lymphoma cell line CD20 is a cell surface 33-37 (depending on the degree of phosphorylation) kDa non-glycosylated surface phosphoprotein expressed on mature and most malignant B cells, but not stem cells or plasma cells (low number of the CD20 has been also detected on Clonality Monoclonal Concentration 1 mg/ml Clone Designation [MEM-97] Isotype IgG1 Phosphate buffered saline (PBS), pH 7.4	Category	Antikörper
Immunogen Raji human Burkitts lymphoma cell line CD20 is a cell surface 33-37 (depending on the degree of phosphorylation) kDa non-glycosylated surface phosphoprotein expressed on mature and most malignant B cells, but not stem cells or plasma cells (low number of the CD20 has been also detected on Clonality Monoclonal Concentration 1 mg/ml Clone Designation [MEM-97] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4	Application	FC, IP
CD20 is a cell surface 33-37 (depending on the degree of phosphorylation) kDa non-glycosylated surface phosphoprotein expressed on mature and most malignant B cells, but not stem cells or plasma cells (low number of the CD20 has been also detected on Clonality Monoclonal Concentration 1 mg/ml Clone Designation [MEM-97] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4	Species Reactivity	Porcine, Human, Bovine
phosphorylation) kDa non-glycosylated surface phosphoprotein expressed on mature and most malignant B cells, but not stem cells or plasma cells (low number of the CD20 has been also detected on Clonality Monoclonal Concentration 1 mg/ml Clone Designation [MEM-97] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4	Immunogen	Raji human Burkitts lymphoma cell line
Concentration 1 mg/ml Clone Designation [MEM-97] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4	Product Description	phosphorylation) kDa non-glycosylated surface phosphoprotein expressed on mature and most malignant B cells, but not stem cells or plasma cells (low number of the CD20 has been also detected
Clone Designation [MEM-97] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4	Clonality	Monoclonal
Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4	Concentration	1 mg/ml
Buffer Phosphate buffered saline (PBS), pH 7.4	Clone Designation	[MEM-97]
	Isotype	lgG1
Storage 2°C to 8°C	Buffer	Phosphate buffered saline (PBS), pH 7.4
	Storage	2°C to 8°C

Antibody Type	Monoclonal Antibody
Application Notes	Flow cytometry: Recommended dilution: 10 µg/ml.