

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Anti-Hu CD40 Purified Azide Free, Clone: [HI40a], Monoclonal EXB-10-416-C100

Article Name Anti-Hu CD40 Purified Azide Free, Clone: [HI40a], Monoclonal  Biozol Catalog Number EXB-10-416-C100  Supplier Catalog Number 10-416-C100  Alternative Catalog Number EXB-10-416-C100  Manufacturer EXBIO  Category Antikörper  Application FC  Species Reactivity Human  Immunogen Human CD40a  CD40 is a costimulatory molecule of the TNF receptor superfamily and is expressed on many cell types, such as B cells, monocytes/macrophages, dendritic cells, endothelial cells, fibroblasts or vascular smooth muscle cells. Interaction of CD40 and its  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [HI40a]  Isotype Mouse IgG1		
Supplier Catalog Number 10-416-C100  Alternative Catalog Number EXB-10-416-C100  Manufacturer EXBIO  Category Antikörper  Application FC  Species Reactivity Human  Immunogen Human CD40a  CD40 is a costimulatory molecule of the TNF receptor superfamily and is expressed on many cell types, such as B cells, monocytes/macrophages, dendritic cells, endothelial cells, fibroblasts or vascular smooth muscle cells. Interaction of CD40 and its  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [HI40a]	Article Name	Anti-Hu CD40 Purified Azide Free, Clone: [HI40a], Monoclonal
Alternative Catalog Number EXB-10-416-C100  Manufacturer EXBIO  Category Antikörper  Application FC  Species Reactivity Human  Immunogen Human CD40a  CD40 is a costimulatory molecule of the TNF receptor superfamily and is expressed on many cell types, such as B cells, monocytes/macrophages, dendritic cells, endothelial cells, fibroblasts or vascular smooth muscle cells. Interaction of CD40 and its  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [HI40a]	Biozol Catalog Number	EXB-10-416-C100
Manufacturer EXBIO  Category Antikörper  Application FC  Species Reactivity Human  Immunogen Human CD40a  CD40 is a costimulatory molecule of the TNF receptor superfamily and is expressed on many cell types, such as B cells, monocytes/macrophages, dendritic cells, endothelial cells, fibroblasts or vascular smooth muscle cells. Interaction of CD40 and its  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [HI40a]	Supplier Catalog Number	10-416-C100
Category Antikörper  Application FC  Species Reactivity Human  Immunogen Human CD40a  CD40 is a costimulatory molecule of the TNF receptor superfamily and is expressed on many cell types, such as B cells, monocytes/macrophages, dendritic cells, endothelial cells, fibroblasts or vascular smooth muscle cells. Interaction of CD40 and its  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [HI40a]	Alternative Catalog Number	EXB-10-416-C100
Application FC  Species Reactivity Human  Immunogen Human CD40a  CD40 is a costimulatory molecule of the TNF receptor superfamily and is expressed on many cell types, such as B cells, monocytes/macrophages, dendritic cells, endothelial cells, fibroblasts or vascular smooth muscle cells. Interaction of CD40 and its  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [HI40a]	Manufacturer	EXBIO
Species Reactivity  Immunogen  Human CD40a  CD40 is a costimulatory molecule of the TNF receptor superfamily and is expressed on many cell types, such as B cells, monocytes/macrophages, dendritic cells, endothelial cells, fibroblasts or vascular smooth muscle cells. Interaction of CD40 and its  Clonality  Monoclonal  Concentration  1 mg/ml  Clone Designation  [HI40a]	Category	Antikörper
Immunogen  Human CD40a  CD40 is a costimulatory molecule of the TNF receptor superfamily and is expressed on many cell types, such as B cells, monocytes/macrophages, dendritic cells, endothelial cells, fibroblasts or vascular smooth muscle cells. Interaction of CD40 and its  Clonality  Monoclonal  Concentration  1 mg/ml  Clone Designation  [HI40a]	Application	FC
CD40 is a costimulatory molecule of the TNF receptor superfamily and is expressed on many cell types, such as B cells, monocytes/macrophages, dendritic cells, endothelial cells, fibroblasts or vascular smooth muscle cells. Interaction of CD40 and its  Clonality  Monoclonal  Concentration  1 mg/ml  Clone Designation  [HI40a]	Species Reactivity	Human
and is expressed on many cell types, such as B cells, monocytes/macrophages, dendritic cells, endothelial cells, fibroblasts or vascular smooth muscle cells. Interaction of CD40 and its  Clonality  Monoclonal  Concentration  1 mg/ml  Clone Designation  [HI40a]	Immunogen	Human CD40a
Concentration 1 mg/ml Clone Designation [HI40a]	Product Description	and is expressed on many cell types, such as B cells, monocytes/macrophages, dendritic cells, endothelial cells, fibroblasts or vascular smooth muscle cells. Interaction of CD40 and
Clone Designation [HI40a]	Clonality	Monoclonal
	Concentration	1 mg/ml
Isotype Mouse IgG1	Clone Designation	[HI40a]
	Isotype	Mouse IgG1
Buffer Phosphate buffered saline (PBS), pH 7.4	Buffer	Phosphate buffered saline (PBS), pH 7.4
Storage 2°C to 8°C	Storage	2°C to 8°C

Target	CD40
Antibody Type	Monoclonal Antibody
Application Notes	Flow cytometry: The purified antibody is effective for indirect immunofluorescence staining of human cells by flow cytometry analysis using 1 $\mu$ g/ml.