

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Hu CD22 Purified Azide Free, Clone: [IS7], Monoclonal EXB-10-506-C025

Article Name Anti-Hu CD22 Purified Azide Free, Clone: [IS7], Monoclonal Biozol Catalog Number EXB-10-506-C025 Supplier Catalog Number 10-506-C025 Alternative Catalog Number EXB-10-506-C025 Manufacturer EXBIO Category Antikörper Application FC Species Reactivity Human Immunogen human cell line Reh CD22, also known as Siglec-2 (sialic acid-binding immunoglobulin-like lectin-2) is a transmembrane glycoprotein binding alpha2,6-linked sialic acid-bearing ligands. Intracellular domain of CD22 recruits protein tyrosine phosphatase SHP-1 through the Clonality Monoclonal Concentration Img/ml Clone Designation [IS7] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4 Storage 2°C to 8°C		
Supplier Catalog Number 10-506-C025 Alternative Catalog Number EXB-10-506-C025 Manufacturer EXBIO Category Antikörper Application FC Species Reactivity Human Immunogen human cell line Reh CD22, also known as Siglec-2 (sialic acid-binding immunoglobulin-like lectin-2) is a transmembrane glycoprotein binding alpha2,6-linked sialic acid-bearing ligands. Intracellular domain of CD22 recruits protein tyrosine phosphatase SHP-1 through the Clonality Monoclonal Concentration 1 mg/ml Clone Designation [IS7] Isotype IgG1 Phosphate buffered saline (PBS), pH 7.4	Article Name	Anti-Hu CD22 Purified Azide Free, Clone: [IS7], Monoclonal
Alternative Catalog Number EXB-10-506-C025 Manufacturer EXBIO Category Antikörper Application FC Species Reactivity Human Immunogen human cell line Reh CD22, also known as Siglec-2 (sialic acid-binding immunoglobulin-like lectin-2) is a transmembrane glycoprotein binding alpha2,6-linked sialic acid-bearing ligands. Intracellular domain of CD22 recruits protein tyrosine phosphatase SHP-1 through the Clonality Monoclonal Concentration I mg/ml Clone Designation [IS7] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4	Biozol Catalog Number	EXB-10-506-C025
Manufacturer EXBIO Category Antikörper Application FC Species Reactivity Human human cell line Reh CD22, also known as Siglec-2 (sialic acid-binding immunoglobulin-like lectin-2) is a transmembrane glycoprotein binding alpha2,6-linked sialic acid-bearing ligands. Intracellular domain of CD22 recruits protein tyrosine phosphatase SHP-1 through the Clonality Monoclonal Concentration I mg/ml Clone Designation [IS7] Isotype IgG1 Phosphate buffered saline (PBS), pH 7.4	Supplier Catalog Number	10-506-C025
Category Antikörper Application FC Species Reactivity Human Immunogen human cell line Reh CD22, also known as Siglec-2 (sialic acid-binding immunoglobulin-like lectin-2) is a transmembrane glycoprotein binding alpha2,6-linked sialic acid-bearing ligands. Intracellular domain of CD22 recruits protein tyrosine phosphatase SHP-1 through the Clonality Monoclonal Concentration 1 mg/ml Clone Designation [IS7] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4	Alternative Catalog Number	EXB-10-506-C025
Application FC Species Reactivity Human Immunogen human cell line Reh CD22, also known as Siglec-2 (sialic acid-binding immunoglobulin-like lectin-2) is a transmembrane glycoprotein binding alpha2,6-linked sialic acid-bearing ligands. Intracellular domain of CD22 recruits protein tyrosine phosphatase SHP-1 through the Clonality Monoclonal Concentration 1 mg/ml Clone Designation [IS7] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4	Manufacturer	EXBIO
Species Reactivity Immunogen human cell line Reh CD22, also known as Siglec-2 (sialic acid-binding immunoglobulin-like lectin-2) is a transmembrane glycoprotein binding alpha2,6-linked sialic acid-bearing ligands. Intracellular domain of CD22 recruits protein tyrosine phosphatase SHP-1 through the Clonality Monoclonal Concentration 1 mg/ml Clone Designation [IS7] Isotype IgG1 Phosphate buffered saline (PBS), pH 7.4	Category	Antikörper
Immunogen human cell line Reh CD22, also known as Siglec-2 (sialic acid-binding immunoglobulin-like lectin-2) is a transmembrane glycoprotein binding alpha2,6-linked sialic acid-bearing ligands. Intracellular domain of CD22 recruits protein tyrosine phosphatase SHP-1 through the Clonality Monoclonal Concentration 1 mg/ml Clone Designation [IS7] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4	Application	FC
Product Description CD22, also known as Siglec-2 (sialic acid-binding immunoglobulin-like lectin-2) is a transmembrane glycoprotein binding alpha2,6-linked sialic acid-bearing ligands. Intracellular domain of CD22 recruits protein tyrosine phosphatase SHP-1 through the Clonality Monoclonal Concentration 1 mg/ml Clone Designation [IS7] Isotype IgG1 Phosphate buffered saline (PBS), pH 7.4	Species Reactivity	Human
Product Description like lectin-2) is a transmembrane glycoprotein binding alpha2,6-linked sialic acid-bearing ligands. Intracellular domain of CD22 recruits protein tyrosine phosphatase SHP-1 through the Clonality Monoclonal Concentration 1 mg/ml Clone Designation [IS7] Isotype IgG1 Phosphate buffered saline (PBS), pH 7.4	Immunogen	human cell line Reh
Concentration 1 mg/ml Clone Designation [IS7] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4	Product Description	like lectin-2) is a transmembrane glycoprotein binding alpha2,6- linked sialic acid-bearing ligands. Intracellular domain of CD22
Clone Designation [IS7] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4	Clonality	Monoclonal
Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4	Concentration	1 mg/ml
Buffer Phosphate buffered saline (PBS), pH 7.4	Clone Designation	[IS7]
	Isotype	lgG1
Storage 2°C to 8°C	Buffer	Phosphate buffered saline (PBS), pH 7.4
	Storage	2°C to 8°C

Antibody Type	Monoclonal Antibody
Application Notes	Flow cytometry: Recommended dilution: 1-4 μ g/ml, Positive control: lysed whole blood, lyse-wash protocol