

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Hu CD150 Purified Azide Free, Clone: [SLAM.4], Monoclonal EXB-10-660-C025

Article Name Anti-Hu CD150 Purified Azide Free, Clone: [SLAM.4], Monoclonal Biozol Catalog Number EXB-10-660-C025 Supplier Catalog Number 10-660-C025 Alternative Catalog Number EXB-10-660-C025 Manufacturer EXBIO Category Antikörper Application FC, ICC, IP Species Reactivity Human Immunogen Human CD150-transfected 300.19 cells CD150, also known as SLAM (signaling lymphocyte activation molecule) is a 70-95 kDa single chain transmembrane phosphoglycoprotein of the CD2 family. Its extracellular part contains eight potential N-glycosylation sites, and the intracellular tail co Clonality Monoclonal Concentration 1 mg/ml Clone Designation [SLAM.4] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4		
Supplier Catalog Number Alternative Catalog Number EXB-10-660-C025 Manufacturer EXBIO Category Antikörper Application FC, ICC, IP Species Reactivity Human Immunogen Human CD150-transfected 300.19 cells CD150, also known as SLAM (signaling lymphocyte activation molecule) is a 70-95 kDa single chain transmembrane phosphoglycoprotein of the CD2 family. Its extracellular part contains eight potential N-glycosylation sites, and the intracellular tail co Clonality Monoclonal Concentration I mg/ml Clone Designation [SLAM.4] Buffer Phosphate buffered saline (PBS), pH 7.4	Article Name	Anti-Hu CD150 Purified Azide Free, Clone: [SLAM.4], Monoclonal
Alternative Catalog Number EXB-10-660-C025 Manufacturer EXBIO Category Antikörper Application FC, ICC, IP Species Reactivity Human Immunogen Human CD150-transfected 300.19 cells CD150, also known as SLAM (signaling lymphocyte activation molecule) is a 70-95 kDa single chain transmembrane phosphoglycoprotein of the CD2 family. Its extracellular part contains eight potential N-glycosylation sites, and the intracellular tail co Clonality Monoclonal Concentration 1 mg/ml Clone Designation [SLAM.4] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4	Biozol Catalog Number	EXB-10-660-C025
Manufacturer EXBIO Category Antikörper Application FC, ICC, IP Species Reactivity Human Human CD150-transfected 300.19 cells CD150, also known as SLAM (signaling lymphocyte activation molecule) is a 70-95 kDa single chain transmembrane phosphoglycoprotein of the CD2 family. Its extracellular part contains eight potential N-glycosylation sites, and the intracellular tail co Clonality Monoclonal Concentration 1 mg/ml Clone Designation [SLAM.4] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4	Supplier Catalog Number	10-660-C025
Category Antikörper Application FC, ICC, IP Species Reactivity Human Immunogen Human CD150-transfected 300.19 cells CD150, also known as SLAM (signaling lymphocyte activation molecule) is a 70-95 kDa single chain transmembrane phosphoglycoprotein of the CD2 family. Its extracellular part contains eight potential N-glycosylation sites, and the intracellular tail co Clonality Monoclonal Concentration 1 mg/ml Clone Designation [SLAM.4] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4	Alternative Catalog Number	EXB-10-660-C025
Application FC, ICC, IP Species Reactivity Human Immunogen Human CD150-transfected 300.19 cells CD150, also known as SLAM (signaling lymphocyte activation molecule) is a 70-95 kDa single chain transmembrane phosphoglycoprotein of the CD2 family. Its extracellular part contains eight potential N-glycosylation sites, and the intracellular tail co Clonality Monoclonal Concentration 1 mg/ml Clone Designation [SLAM.4] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4	Manufacturer	EXBIO
Species Reactivity Human Human CD150-transfected 300.19 cells CD150, also known as SLAM (signaling lymphocyte activation molecule) is a 70-95 kDa single chain transmembrane phosphoglycoprotein of the CD2 family. Its extracellular part contains eight potential N-glycosylation sites, and the intracellular tail co Clonality Monoclonal Concentration 1 mg/ml Clone Designation [SLAM.4] Isotype IgG1 Phosphate buffered saline (PBS), pH 7.4	Category	Antikörper
Immunogen Human CD150-transfected 300.19 cells CD150, also known as SLAM (signaling lymphocyte activation molecule) is a 70-95 kDa single chain transmembrane phosphoglycoprotein of the CD2 family. Its extracellular part contains eight potential N-glycosylation sites, and the intracellular tail co Clonality Monoclonal Concentration 1 mg/ml Clone Designation [SLAM.4] Isotype IgG1 Phosphate buffered saline (PBS), pH 7.4	Application	FC, ICC, IP
CD150, also known as SLAM (signaling lymphocyte activation molecule) is a 70-95 kDa single chain transmembrane phosphoglycoprotein of the CD2 family. Its extracellular part contains eight potential N-glycosylation sites, and the intracellular tail co Clonality Monoclonal Concentration 1 mg/ml Clone Designation [SLAM.4] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4	Species Reactivity	Human
molecule) is a 70-95 kDa single chain transmembrane phosphoglycoprotein of the CD2 family. Its extracellular part contains eight potential N-glycosylation sites, and the intracellular tail co Clonality Monoclonal Concentration 1 mg/ml Clone Designation [SLAM.4] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4	Immunogen	Human CD150-transfected 300.19 cells
Concentration 1 mg/ml Clone Designation [SLAM.4] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4	Product Description	molecule) is a 70-95 kDa single chain transmembrane phosphoglycoprotein of the CD2 family. Its extracellular part contains eight potential N-glycosylation sites, and the intracellular
Clone Designation [SLAM.4] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4	Clonality	Monoclonal
Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4	Concentration	1 mg/ml
Buffer Phosphate buffered saline (PBS), pH 7.4	Clone Designation	[SLAM.4]
	Isotype	IgG1
	Buffer	Phosphate buffered saline (PBS), pH 7.4
Storage 2°C to 8°C	Storage	2°C to 8°C

Antibody Type	Monoclonal Antibody
Application Notes	Flow cytometry: Recommended dilution: 1-3 μg/ml.