

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Hu CD164 Purified, Clone: [67D2], Monoclonal EXB-11-135-C025

| Article Name Anti-Hu CD164 Purified, Clone: [67D2], Monoclonal Biozol Catalog Number EXB-11-135-C025 Supplier Catalog Number 11-135-C025 Alternative Catalog Number EXB-11-135-C025 Manufacturer EXBIO Category Antikörper Application WB, FC, ICC, IP Species Reactivity Human Immunogen Breast tumor cell line T-47D CD164, also known as endolyn, is a type I transmembrane protein with heavily glycosylated extracellular part containing sialic acid ar glycosaminoglycan residues. CD164 plays both adhesive and antiadhesive role and serves as a potent negative regula Clonality Monoclonal |
|---|
| Supplier Catalog Number Alternative Catalog Number EXB-11-135-C025 Manufacturer EXBIO Category Antikörper Application WB, FC, ICC, IP Species Reactivity Human Immunogen Breast tumor cell line T-47D CD164, also known as endolyn, is a type I transmembrane protein with heavily glycosylated extracellular part containing sialic acid ar glycosaminoglycan residues. CD164 plays both adhesive and antiadhesive role and serves as a potent negative regula |
| Alternative Catalog Number EXB-11-135-C025 Manufacturer EXBIO Category Antikörper Application WB, FC, ICC, IP Species Reactivity Human Immunogen Breast tumor cell line T-47D CD164, also known as endolyn, is a type I transmembrane protein with heavily glycosylated extracellular part containing sialic acid ar glycosaminoglycan residues. CD164 plays both adhesive and antiadhesive role and serves as a potent negative regula |
| Manufacturer EXBIO Category Antikörper Application WB, FC, ICC, IP Species Reactivity Human Immunogen Breast tumor cell line T-47D CD164, also known as endolyn, is a type I transmembrane protein with heavily glycosylated extracellular part containing sialic acid ar glycosaminoglycan residues. CD164 plays both adhesive and antiadhesive role and serves as a potent negative regula |
| Category Antikörper Application WB, FC, ICC, IP Species Reactivity Human Immunogen Breast tumor cell line T-47D CD164, also known as endolyn, is a type I transmembrane protein with heavily glycosylated extracellular part containing sialic acid ar glycosaminoglycan residues. CD164 plays both adhesive and antiadhesive role and serves as a potent negative regula |
| Application WB, FC, ICC, IP Species Reactivity Human Breast tumor cell line T-47D CD164, also known as endolyn, is a type I transmembrane protein with heavily glycosylated extracellular part containing sialic acid ar glycosaminoglycan residues. CD164 plays both adhesive and antiadhesive role and serves as a potent negative regula |
| Species Reactivity Human Breast tumor cell line T-47D CD164, also known as endolyn, is a type I transmembrane protein with heavily glycosylated extracellular part containing sialic acid ar glycosaminoglycan residues. CD164 plays both adhesive and antiadhesive role and serves as a potent negative regula |
| Immunogen Breast tumor cell line T-47D CD164, also known as endolyn, is a type I transmembrane protein with heavily glycosylated extracellular part containing sialic acid ar glycosaminoglycan residues. CD164 plays both adhesive and antiadhesive role and serves as a potent negative regula |
| Product Description Product Description CD164, also known as endolyn, is a type I transmembrane protein with heavily glycosylated extracellular part containing sialic acid ar glycosaminoglycan residues. CD164 plays both adhesive and antiadhesive role and serves as a potent negative regula |
| Product Description with heavily glycosylated extracellular part containing sialic acid an glycosaminoglycan residues. CD164 plays both adhesive and antiadhesive role and serves as a potent negative regula |
| Clonality Monoclonal |
| indictional indictional |
| Concentration 1 mg/ml |
| Clone Designation [67D2] |
| Isotype IgG1 |
| Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide |
| Storage 2°C to 8°C |

| Antibody Type | Monoclonal Antibody |
|-------------------|--|
| Application Notes | Flow cytometry: Recommended dilution: 1-4 μg/ml. |