

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Anti-Hu CD157 Purified, Clone: [SY11B5], Monoclonal EXB-11-136-C025

Article Name Anti-Hu CD157 Purified, Clone: [SY11B5], Monoclonal  Biozol Catalog Number EXB-11-136-C025  Supplier Catalog Number 11-136-C025  Alternative Catalog Number EXB-11-136-C025  Manufacturer EXBIO  Category Antikörper  Application WB, FC, IHC-Fr, IP  Species Reactivity Primate, Human  Immunogen Human CD157  CD157 (cADPr hydrolase 2) is a GPI-anchored ectoenzyme possessing ADP-ribosyl cyclase and cyclic ADP-ribose hydrolase activity. It uses NAD and cADP-ribose as substrates. CD157 is expressed on granulocytes, monocytes, macrophages, follicular dendriti  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [SY11B5]  Isotype IgG1, kappa  Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide  Storage 2°C to 8°C		
Supplier Catalog Number 11-136-C025  Alternative Catalog Number EXB-11-136-C025  Manufacturer EXBIO  Category Antikörper  Application WB, FC, IHC-Fr, IP  Species Reactivity Primate, Human  Immunogen Human CD157  CD157 (cADPr hydrolase 2) is a GPI-anchored ectoenzyme possessing ADP-ribosyl cyclase and cyclic ADP-ribose hydrolase activity. It uses NAD and cADP-ribose as substrates. CD157 is expressed on granulocytes, monocytes, macrophages, follicular dendriti  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [SY11B5]  Isotype IgG1, kappa  Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Article Name	Anti-Hu CD157 Purified, Clone: [SY11B5], Monoclonal
Alternative Catalog Number  EXB-11-136-C025  Manufacturer  EXBIO  Category  Antikörper  Application  WB, FC, IHC-Fr, IP  Species Reactivity  Primate, Human  Immunogen  Human CD157  CD157 (cADPr hydrolase 2) is a GPI-anchored ectoenzyme possessing ADP-ribosyl cyclase and cyclic ADP-ribose hydrolase activity. It uses NAD and cADP-ribose as substrates. CD157 is expressed on granulocytes, monocytes, macrophages, follicular dendriti  Clonality  Monoclonal  Concentration  1 mg/ml  Clone Designation  [SY11B5]  Isotype  IgG1, kappa  Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Biozol Catalog Number	EXB-11-136-C025
Manufacturer  EXBIO  Category  Antikörper  Application  WB, FC, IHC-Fr, IP  Species Reactivity  Primate, Human  Immunogen  Human CD157  CD157 (cADPr hydrolase 2) is a GPI-anchored ectoenzyme possessing ADP-ribosyl cyclase and cyclic ADP-ribose hydrolase activity. It uses NAD and cADP-ribose as substrates. CD157 is expressed on granulocytes, monocytes, macrophages, follicular dendriti  Clonality  Monoclonal  Concentration  1 mg/ml  Clone Designation  [SY11B5]  Isotype  IgG1, kappa  Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Supplier Catalog Number	11-136-C025
Category Antikörper  Application WB, FC, IHC-Fr, IP  Species Reactivity Primate, Human  Immunogen Human CD157  CD157 (cADPr hydrolase 2) is a GPI-anchored ectoenzyme possessing ADP-ribosyl cyclase and cyclic ADP-ribose hydrolase activity. It uses NAD and cADP-ribose as substrates. CD157 is expressed on granulocytes, monocytes, macrophages, follicular dendriti  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [SY11B5]  Isotype IgG1, kappa  Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Alternative Catalog Number	EXB-11-136-C025
Application  WB, FC, IHC-Fr, IP  Species Reactivity  Primate, Human  Human CD157  CD157 (cADPr hydrolase 2) is a GPI-anchored ectoenzyme possessing ADP-ribosyl cyclase and cyclic ADP-ribose hydrolase activity. It uses NAD and cADP-ribose as substrates. CD157 is expressed on granulocytes, monocytes, macrophages, follicular dendriti  Clonality  Monoclonal  Concentration  1 mg/ml  Clone Designation  [SY11B5]  Isotype  IgG1, kappa  Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Manufacturer	EXBIO
Species Reactivity Primate, Human  Human CD157  CD157 (cADPr hydrolase 2) is a GPI-anchored ectoenzyme possessing ADP-ribosyl cyclase and cyclic ADP-ribose hydrolase activity. It uses NAD and cADP-ribose as substrates. CD157 is expressed on granulocytes, monocytes, macrophages, follicular dendriti  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [SY11B5]  Isotype IgG1, kappa  Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Category	Antikörper
Immunogen       Human CD157         CD157 (cADPr hydrolase 2) is a GPI-anchored ectoenzyme possessing ADP-ribosyl cyclase and cyclic ADP-ribose hydrolase activity. It uses NAD and cADP-ribose as substrates. CD157 is expressed on granulocytes, monocytes, macrophages, follicular dendriti         Clonality       Monoclonal         Concentration       1 mg/ml         Clone Designation       [SY11B5]         Isotype       IgG1, kappa         Buffer       Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Application	WB, FC, IHC-Fr, IP
CD157 (cADPr hydrolase 2) is a GPI-anchored ectoenzyme possessing ADP-ribosyl cyclase and cyclic ADP-ribose hydrolase activity. It uses NAD and cADP-ribose as substrates. CD157 is expressed on granulocytes, monocytes, macrophages, follicular dendriti  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [SY11B5]  Isotype IgG1, kappa  Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Species Reactivity	Primate, Human
Product Description  possessing ADP-ribosyl cyclase and cyclic ADP-ribose hydrolase activity. It uses NAD and cADP-ribose as substrates. CD157 is expressed on granulocytes, monocytes, macrophages, follicular dendriti  Clonality  Monoclonal  Concentration  1 mg/ml  Clone Designation  [SY11B5]  Isotype  IgG1, kappa  Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Immunogen	Human CD157
Concentration 1 mg/ml Clone Designation [SY11B5] Isotype IgG1, kappa  Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Product Description	possessing ADP-ribosyl cyclase and cyclic ADP-ribose hydrolase activity. It uses NAD and cADP-ribose as substrates. CD157 is expressed on granulocytes, monocytes, macrophages, follicular
Clone Designation [SY11B5]  Isotype IgG1, kappa  Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Clonality	Monoclonal
Isotype IgG1, kappa  Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Concentration	1 mg/ml
Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Clone Designation	[SY11B5]
	Isotype	IgG1, kappa
Storage 2°C to 8°C	Buffer	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
	Storage	2°C to 8°C

Antibody Type	Monoclonal Antibody
Application Notes	Western blotting: Non-reducing conditions. Immunohistochemistry: Recommended dilution: 5-10 $\mu$ g/ml. Flow cytometry: Recommended dilution: 1-12 $\mu$ g/ml