

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | Fax: +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Hu CD57 Purified, Clone: [TB01], Monoclonal EXB-11-158-C025

Article NameAnti-Hu CD57 Purified, Clone: [TB01], MonoclonalBiozol Catalog NumberEXB-11-158-C025Supplier Catalog Number11-158-C025Alternative Catalog NumberEXB-11-158-C025ManufacturerEXBIOCategoryAntikörperApplicationFC, IHC-P, ICC, WB, IHC-FrSpecies ReactivityHumanImmunogenA pool of neuroblastoma cell linesProduct DescriptionCD57, also known as HNK1 or Leu7, is a sulphated trisaccharide (3-O-sulfoglucuronic acid beta1-3 Gal beta1-4 GlcNAc) attached to several glycoproteins, including CD56, myelin glycoprotein PO, and neural cell adhesion molecule L1, as well as on glycolClonalityMonoclonalConcentration1 mg/mlClone Designation[TB01]IsotypeIgMBufferTris buffered saline (TBS), pH 8.0, 15 mM sodium azideStorage2°C to 8°C		
Supplier Catalog Number 11-158-C025 Alternative Catalog Number EXB-11-158-C025 Manufacturer EXBIO Category Antikörper Application FC, IHC-P, ICC, WB, IHC-Fr Species Reactivity Human Immunogen A pool of neuroblastoma cell lines CD57, also known as HNK1 or Leu7, is a sulphated trisaccharide (3-O-sulfoglucuronic acid beta1-3 Gal beta1-4 GlcNAc) attached to several glycoproteins, including CD56, myelin glycoprotein PO, and neural cell adhesion molecule L1, as well as on glycol Clonality Monoclonal Concentration 1 mg/ml Clone Designation [TB01] Isotype IgM Buffer Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Article Name	Anti-Hu CD57 Purified, Clone: [TB01], Monoclonal
Alternative Catalog Number EXB-11-158-C025 Manufacturer EXBIO Category Antikörper Application FC, IHC-P, ICC, WB, IHC-Fr Species Reactivity Human Immunogen A pool of neuroblastoma cell lines CD57, also known as HNK1 or Leu7, is a sulphated trisaccharide (3-O-sulfoglucuronic acid beta1-3 Gal beta1-4 GlcNAc) attached to several glycoproteins, including CD56, myelin glycoprotein PO, and neural cell adhesion molecule L1, as well as on glycol Clonality Monoclonal Concentration 1 mg/ml Clone Designation [TB01] Isotype IgM Buffer Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Biozol Catalog Number	EXB-11-158-C025
Manufacturer EXBIO Category Antikörper Application FC, IHC-P, ICC, WB, IHC-Fr Species Reactivity Human Immunogen A pool of neuroblastoma cell lines CD57, also known as HNK1 or Leu7, is a sulphated trisaccharide (3-O-sulfoglucuronic acid beta1-3 Gal beta1-4 GlcNAc) attached to several glycoproteins, including CD56, myelin glycoprotein PO, and neural cell adhesion molecule L1, as well as on glycol Clonality Monoclonal Concentration I mg/ml Clone Designation [TB01] Isotype IgM Buffer Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Supplier Catalog Number	11-158-C025
Category Application FC, IHC-P, ICC, WB, IHC-Fr Species Reactivity Human A pool of neuroblastoma cell lines CD57, also known as HNK1 or Leu7, is a sulphated trisaccharide (3-O-sulfoglucuronic acid beta1-3 Gal beta1-4 GlcNAc) attached to several glycoproteins, including CD56, myelin glycoprotein PO, and neural cell adhesion molecule L1, as well as on glycol Clonality Monoclonal Concentration 1 mg/ml Clone Designation [TB01] Isotype IgM Buffer Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Alternative Catalog Number	EXB-11-158-C025
Application FC, IHC-P, ICC, WB, IHC-Fr Species Reactivity Human Immunogen A pool of neuroblastoma cell lines CD57, also known as HNK1 or Leu7, is a sulphated trisaccharide (3-O-sulfoglucuronic acid beta1-3 Gal beta1-4 GlcNAc) attached to several glycoproteins, including CD56, myelin glycoprotein PO, and neural cell adhesion molecule L1, as well as on glycol Clonality Monoclonal Concentration 1 mg/ml Clone Designation [TB01] Isotype IgM Buffer Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Manufacturer	EXBIO
Species Reactivity Immunogen A pool of neuroblastoma cell lines CD57, also known as HNK1 or Leu7, is a sulphated trisaccharide (3-O-sulfoglucuronic acid beta1-3 Gal beta1-4 GlcNAc) attached to several glycoproteins, including CD56, myelin glycoprotein PO, and neural cell adhesion molecule L1, as well as on glycol Clonality Monoclonal Concentration 1 mg/ml Clone Designation [TB01] Isotype IgM Buffer Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Category	Antikörper
Immunogen A pool of neuroblastoma cell lines Product Description CD57, also known as HNK1 or Leu7, is a sulphated trisaccharide (3-O-sulfoglucuronic acid beta1-3 Gal beta1-4 GlcNAc) attached to several glycoproteins, including CD56, myelin glycoprotein PO, and neural cell adhesion molecule L1, as well as on glycol Clonality Monoclonal Concentration 1 mg/ml Clone Designation [TB01] Isotype IgM Buffer Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Application	FC, IHC-P, ICC, WB, IHC-Fr
Product Description CD57, also known as HNK1 or Leu7, is a sulphated trisaccharide (3-O-sulfoglucuronic acid beta1-3 Gal beta1-4 GlcNAc) attached to several glycoproteins, including CD56, myelin glycoprotein PO, and neural cell adhesion molecule L1, as well as on glycol Clonality Monoclonal Concentration 1 mg/ml Clone Designation [TB01] Isotype IgM Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Species Reactivity	Human
Product Description sulfoglucuronic acid beta1-3 Gal beta1-4 GlcNAc) attached to several glycoproteins, including CD56, myelin glycoprotein PO, and neural cell adhesion molecule L1, as well as on glycol Clonality Monoclonal Concentration 1 mg/ml Clone Designation [TB01] Isotype IgM Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Immunogen	A pool of neuroblastoma cell lines
Concentration 1 mg/ml Clone Designation [TB01] Isotype IgM Buffer Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Product Description	sulfoglucuronic acid beta1-3 Gal beta1-4 GlcNAc) attached to several glycoproteins, including CD56, myelin glycoprotein PO, and neural
Clone Designation [TB01] Isotype IgM Buffer Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Clonality	Monoclonal
Isotype IgM Buffer Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Concentration	1 mg/ml
Buffer Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Clone Designation	[TB01]
The same ca same (125), pit die, 25 mil soaiam aziac	Isotype	IgM
Storage 2°C to 8°C	Buffer	Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide
	Storage	2°C to 8°C

Antibody Type	Monoclonal Antibody
Application Notes	Flow cytometry: Recommended dilution: 1-4 μg/ml. Immunohistochemistry: Recommended dilution: 5-10 μg/ml.