

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | Fax: +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Anti-Hu CD11b Purified, Clone: [MEM-174], Monoclonal EXB-11-211-C025

Article Name Anti-Hu CD11b Purified, Clone: [MEM-174], Monoclonal  Biozol Catalog Number EXB-11-211-C025  Supplier Catalog Number 11-211-C025  Alternative Catalog Number EXB-11-211-C025  Manufacturer EXBIO  Category Antikörper  Application FC, IP  Species Reactivity Human  Immunogen Human granulocytes  CD11b (integrin alphaM subunit) is a 165-170 kDa type I transmembrane glycoprotein that non-covalently associates with integrin beta2 subunit (CD18), expression of the CD11b chain on the cell surface requires the presence of the CD18 antigen. CD11b/C  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [MEM-174]  Isotype IgG2a  Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide		
Supplier Catalog Number 11-211-C025  Alternative Catalog Number EXB-11-211-C025  Manufacturer EXBIO  Category Antikörper  Application FC, IP  Species Reactivity Human  Immunogen Human granulocytes  CD11b (integrin alphaM subunit) is a 165-170 kDa type I transmembrane glycoprotein that non-covalently associates with integrin beta2 subunit (CD18), expression of the CD11b chain on the cell surface requires the presence of the CD18 antigen. CD11b/C  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [MEM-174]  Isotype IgG2a	Article Name	Anti-Hu CD11b Purified, Clone: [MEM-174], Monoclonal
Alternative Catalog Number EXB-11-211-C025  Manufacturer EXBIO  Category Antikörper  Application FC, IP  Species Reactivity Human  Immunogen Human granulocytes  CD11b (integrin alphaM subunit) is a 165-170 kDa type I transmembrane glycoprotein that non-covalently associates with integrin beta2 subunit (CD18), expression of the CD11b chain on the cell surface requires the presence of the CD18 antigen. CD11b/C  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [MEM-174]  Isotype IgG2a	Biozol Catalog Number	EXB-11-211-C025
Manufacturer  EXBIO  Category  Antikörper  Application  FC, IP  Species Reactivity  Human  Immunogen  Human granulocytes  CD11b (integrin alphaM subunit) is a 165-170 kDa type I transmembrane glycoprotein that non-covalently associates with integrin beta2 subunit (CD18), expression of the CD11b chain on the cell surface requires the presence of the CD18 antigen. CD11b/C  Clonality  Monoclonal  Concentration  1 mg/ml  Clone Designation  [MEM-174]  Isotype  IgG2a	Supplier Catalog Number	11-211-C025
Category Antikörper  Application FC, IP  Species Reactivity Human  Immunogen Human granulocytes  CD11b (integrin alphaM subunit) is a 165-170 kDa type I transmembrane glycoprotein that non-covalently associates with integrin beta2 subunit (CD18), expression of the CD11b chain on the cell surface requires the presence of the CD18 antigen. CD11b/C  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [MEM-174]  Isotype IgG2a	Alternative Catalog Number	EXB-11-211-C025
Application FC, IP  Species Reactivity Human  Immunogen Human granulocytes  CD11b (integrin alphaM subunit) is a 165-170 kDa type I transmembrane glycoprotein that non-covalently associates with integrin beta2 subunit (CD18), expression of the CD11b chain on the cell surface requires the presence of the CD18 antigen. CD11b/C  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [MEM-174]  Isotype IgG2a	Manufacturer	EXBIO
Species Reactivity  Immunogen  Human granulocytes  CD11b (integrin alphaM subunit) is a 165-170 kDa type I transmembrane glycoprotein that non-covalently associates with integrin beta2 subunit (CD18), expression of the CD11b chain on the cell surface requires the presence of the CD18 antigen. CD11b/C  Clonality  Monoclonal  Concentration  1 mg/ml  Clone Designation  [MEM-174]  Isotype  IgG2a	Category	Antikörper
Immunogen       Human granulocytes         CD11b (integrin alphaM subunit) is a 165-170 kDa type I transmembrane glycoprotein that non-covalently associates with integrin beta2 subunit (CD18), expression of the CD11b chain on the cell surface requires the presence of the CD18 antigen. CD11b/C         Clonality       Monoclonal         Concentration       1 mg/ml         Clone Designation       [MEM-174]         Isotype       IgG2a	Application	FC, IP
Product Description  CD11b (integrin alphaM subunit) is a 165-170 kDa type I transmembrane glycoprotein that non-covalently associates with integrin beta2 subunit (CD18), expression of the CD11b chain on the cell surface requires the presence of the CD18 antigen. CD11b/C  Clonality  Monoclonal  Concentration  1 mg/ml  Clone Designation  [MEM-174]  Isotype  IgG2a	Species Reactivity	Human
Product Description transmembrane glycoprotein that non-covalently associates with integrin beta2 subunit (CD18), expression of the CD11b chain on the cell surface requires the presence of the CD18 antigen. CD11b/C  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [MEM-174]  Isotype IgG2a	Immunogen	Human granulocytes
Concentration 1 mg/ml Clone Designation [MEM-174] Isotype IgG2a	Product Description	transmembrane glycoprotein that non-covalently associates with integrin beta2 subunit (CD18), expression of the CD11b chain on the
Clone Designation [MEM-174]  Isotype IgG2a	Clonality	Monoclonal
Isotype IgG2a	Concentration	1 mg/ml
	Clone Designation	[MEM-174]
Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Isotype	IgG2a
	Buffer	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Storage 2°C to 8°C	Storage	2°C to 8°C

Antibody Type	Monoclonal Antibody
Application Notes	Flow cytometry: Recommended dilution: 1 $\mu$ g/ml, positive control: peripheral blood lymphocytes. Immunoprecipitation: Positive control: granulocytes, CD11b/CD18 transfectants. Immunoprecipitation with the antibody MEM-174 was successfuly performed