

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Hu CD11b Purified, Clone: [MEM-174], Monoclonal EXB-11-211-C100

Article Name Anti-Hu CD11b Purified, Clone: [MEM-174], Monoclonal Biozol Catalog Number EXB-11-211-C100 Supplier Catalog Number 11-211-C100 Alternative Catalog Number EXB-11-211-C100 Manufacturer EXBIO Category Antikörper Application FC, IP Species Reactivity Human Immunogen Human granulocytes CD11b (integrin alphaM subunit) is a 165-170 kDa type I transmembrane glycoprotein that non-covalently associates with integrin beta2 subunit (CD18), expression of the CD11b chain on the cell surface requires the presence of the CD18 antigen. CD11b/C Clonality Monoclonal Concentration I mg/ml Clone Designation [MEM-174] Isotype Mouse IgG2a Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide Storage 2°C to 8°C		
Supplier Catalog Number Alternative Catalog Number EXB-11-211-C100 Manufacturer EXBIO Category Antikörper Application FC, IP Species Reactivity Human Immunogen Human granulocytes CD11b (integrin alphaM subunit) is a 165-170 kDa type I transmembrane glycoprotein that non-covalently associates with integrin beta2 subunit (CD18), expression of the CD11b chain on the cell surface requires the presence of the CD18 antigen. CD11b/C Clonality Monoclonal Concentration 1 mg/ml Clone Designation [MEM-174] Isotype Mouse IgG2a Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Article Name	Anti-Hu CD11b Purified, Clone: [MEM-174], Monoclonal
Alternative Catalog Number EXB-11-211-C100 Manufacturer EXBIO Category Antikörper Application FC, IP Species Reactivity Human Immunogen Human granulocytes CD11b (integrin alphaM subunit) is a 165-170 kDa type I transmembrane glycoprotein that non-covalently associates with integrin beta2 subunit (CD18), expression of the CD11b chain on the cell surface requires the presence of the CD18 antigen. CD11b/C Clonality Monoclonal Concentration 1 mg/ml Clone Designation [MEM-174] Isotype Mouse IgG2a Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Biozol Catalog Number	EXB-11-211-C100
Manufacturer EXBIO Category Antikörper Application FC, IP Species Reactivity Human Human granulocytes CD11b (integrin alphaM subunit) is a 165-170 kDa type I transmembrane glycoprotein that non-covalently associates with integrin beta2 subunit (CD18), expression of the CD11b chain on the cell surface requires the presence of the CD18 antigen. CD11b/C Clonality Monoclonal Concentration 1 mg/ml Clone Designation [MEM-174] Isotype Mouse IgG2a Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Supplier Catalog Number	11-211-C100
Category Antikörper Application FC, IP Species Reactivity Human Immunogen Human granulocytes CD11b (integrin alphaM subunit) is a 165-170 kDa type I transmembrane glycoprotein that non-covalently associates with integrin beta2 subunit (CD18), expression of the CD11b chain on the cell surface requires the presence of the CD18 antigen. CD11b/C Clonality Monoclonal Concentration 1 mg/ml Clone Designation [MEM-174] Isotype Mouse IgG2a Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Alternative Catalog Number	EXB-11-211-C100
Application FC, IP Species Reactivity Human Immunogen Human granulocytes CD11b (integrin alphaM subunit) is a 165-170 kDa type I transmembrane glycoprotein that non-covalently associates with integrin beta2 subunit (CD18), expression of the CD11b chain on the cell surface requires the presence of the CD18 antigen. CD11b/C Clonality Monoclonal Concentration 1 mg/ml Clone Designation [MEM-174] Isotype Mouse IgG2a Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Manufacturer	EXBIO
Species Reactivity Immunogen Human granulocytes CD11b (integrin alphaM subunit) is a 165-170 kDa type I transmembrane glycoprotein that non-covalently associates with integrin beta2 subunit (CD18), expression of the CD11b chain on the cell surface requires the presence of the CD18 antigen. CD11b/C Clonality Monoclonal Concentration I mg/ml Clone Designation [MEM-174] Isotype Mouse IgG2a Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Category	Antikörper
Immunogen Human granulocytes CD11b (integrin alphaM subunit) is a 165-170 kDa type I transmembrane glycoprotein that non-covalently associates with integrin beta2 subunit (CD18), expression of the CD11b chain on the cell surface requires the presence of the CD18 antigen. CD11b/C Clonality Monoclonal Concentration 1 mg/ml Clone Designation [MEM-174] Isotype Mouse IgG2a Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Application	FC, IP
Product Description CD11b (integrin alphaM subunit) is a 165-170 kDa type I transmembrane glycoprotein that non-covalently associates with integrin beta2 subunit (CD18), expression of the CD11b chain on the cell surface requires the presence of the CD18 antigen. CD11b/C Clonality Monoclonal Concentration 1 mg/ml Clone Designation [MEM-174] Isotype Mouse IgG2a Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Species Reactivity	Human
Product Description transmembrane glycoprotein that non-covalently associates with integrin beta2 subunit (CD18), expression of the CD11b chain on the cell surface requires the presence of the CD18 antigen. CD11b/C Clonality Monoclonal Concentration 1 mg/ml Clone Designation [MEM-174] Isotype Mouse IgG2a Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Immunogen	Human granulocytes
Concentration 1 mg/ml Clone Designation [MEM-174] Isotype Mouse IgG2a Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Product Description	transmembrane glycoprotein that non-covalently associates with integrin beta2 subunit (CD18), expression of the CD11b chain on the
Clone Designation [MEM-174] Isotype Mouse IgG2a Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Clonality	Monoclonal
Isotype Mouse IgG2a Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Concentration	1 mg/ml
Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Clone Designation	[MEM-174]
Thospitate same (FBS), pit 7.4, 13 mm sociam azide	Isotype	Mouse IgG2a
Storage 2°C to 8°C	Buffer	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
	Storage	2°C to 8°C

Target	CD11b
Antibody Type	Monoclonal Antibody
Application Notes	Flow cytometry: Recommended dilution: 1 μ g/ml, positive control: peripheral blood lymphocytes. Immunoprecipitation: Positive control: granulocytes, CD11b/CD18 transfectants. Immunoprecipitation with the antibody MEM-174 was successfuly performed using immuno-affinity sorbents (minicolumns). The final immunosorbent should contain 1-5 mg of antibody/1 ml of sorbent gel (e.g. agarose beads). It is recommended to stimulate positive control material by LPS 1-2 days before to achieve better expression of target antigen.