

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Anti-Vimentin Purified, Clone: [VI-01], Monoclonal EXB-11-254-C025

Anti-Vimentin Purified, Clone: [VI-01], Monoclonal
EXB-11-254-C025
11-254-C025
EXB-11-254-C025
EXBIO
Antikörper
WB, IHC-P, ICC
Mammal
Pellet of porcine brain cold stable proteins after depolymerization of microtubules.
Vimentin (57 kDa) is the most ubiquituos intermediate filament protein and the first to be expressed during cell differentiation. All primitive cell types express vimentin but in most non-mesenchymal cells it is replaced by other intermediate filamen
Monoclonal
1 mg/ml
[VI-01]
IgM
Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide
2°C to 8°C

Antibody Type	Monoclonal Antibody
Application Notes	Immunocytochemistry: Staining technique: (a) fix cells for 10 min in methanol at -20C and for 6 min in acetone at -20C, (b) fix cells directly in methanol for 10 min at -20C or in acetone for 10 min at -20C. Positive control: 3T3 murine Swiss albino