

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-LCK Purified, Clone: [LCK-01], Monoclonal EXB-11-269-C025

Article Name Anti-LCK Purified, Clone: [LCK-01], Monoclonal Biozol Catalog Number EXB-11-269-C025 Supplier Catalog Number 11-269-C025 Alternative Catalog Number EXB-11-269-C025 Manufacturer EXBIO Category Antikörper Application WB, IP, FC, ICC Species Reactivity Human Immunogen Peptide corresponding to amino acids 22-36 in the sequence of human Lck. Lck is a Src-family tyrosine kinase, which is essential for signaling through the T-cell receptor (TCR) complex. Upon TCR triggering, Lck phosphorylates the ITAM motives in its zeta subunits, establishing binding sites for the SH2 domains of the tyro Clonality Monoclonal Concentration 1 mg/ml Clone Designation [LCK-01] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide		
Supplier Catalog Number Alternative Catalog Number EXB-11-269-C025 Manufacturer EXBIO Category Antikörper Application WB, IP, FC, ICC Species Reactivity Human Peptide corresponding to amino acids 22-36 in the sequence of human Lck. Lck is a Src-family tyrosine kinase, which is essential for signaling through the T-cell receptor (TCR) complex. Upon TCR triggering, Lck phosphorylates the ITAM motives in its zeta subunits, establishing binding sites for the SH2 domains of the tyro Clonality Monoclonal Concentration I mg/ml Clone Designation IgG1	Article Name	Anti-LCK Purified, Clone: [LCK-01], Monoclonal
Alternative Catalog Number EXB-11-269-C025 Manufacturer EXBIO Category Antikörper Application WB, IP, FC, ICC Species Reactivity Human Peptide corresponding to amino acids 22-36 in the sequence of human Lck. Lck is a Src-family tyrosine kinase, which is essential for signaling through the T-cell receptor (TCR) complex. Upon TCR triggering, Lck phosphorylates the ITAM motives in its zeta subunits, establishing binding sites for the SH2 domains of the tyro Clonality Monoclonal Concentration I mg/ml Clone Designation [LCK-01] Isotype IgG1	Biozol Catalog Number	EXB-11-269-C025
Manufacturer EXBIO Category Antikörper Application WB, IP, FC, ICC Species Reactivity Human Peptide corresponding to amino acids 22-36 in the sequence of human Lck. Lck is a Src-family tyrosine kinase, which is essential for signaling through the T-cell receptor (TCR) complex. Upon TCR triggering, Lck phosphorylates the ITAM motives in its zeta subunits, establishing binding sites for the SH2 domains of the tyro Clonality Monoclonal Concentration 1 mg/ml Clone Designation [LCK-01] Isotype	Supplier Catalog Number	11-269-C025
Category Antikörper Application WB, IP, FC, ICC Species Reactivity Human Peptide corresponding to amino acids 22-36 in the sequence of human Lck. Lck is a Src-family tyrosine kinase, which is essential for signaling through the T-cell receptor (TCR) complex. Upon TCR triggering, Lck phosphorylates the ITAM motives in its zeta subunits, establishing binding sites for the SH2 domains of the tyro Clonality Monoclonal Concentration 1 mg/ml Clone Designation [LCK-01] Isotype IgG1	Alternative Catalog Number	EXB-11-269-C025
Application WB, IP, FC, ICC Species Reactivity Human Peptide corresponding to amino acids 22-36 in the sequence of human Lck. Lck is a Src-family tyrosine kinase, which is essential for signaling through the T-cell receptor (TCR) complex. Upon TCR triggering, Lck phosphorylates the ITAM motives in its zeta subunits, establishing binding sites for the SH2 domains of the tyro Clonality Monoclonal Concentration I mg/ml Clone Designation [LCK-01] Isotype IgG1	Manufacturer	EXBIO
Species Reactivity	Category	Antikörper
Immunogen Peptide corresponding to amino acids 22-36 in the sequence of human Lck. Product Description Lck is a Src-family tyrosine kinase, which is essential for signaling through the T-cell receptor (TCR) complex. Upon TCR triggering, Lck phosphorylates the ITAM motives in its zeta subunits, establishing binding sites for the SH2 domains of the tyro Clonality Monoclonal Concentration 1 mg/ml Clone Designation [LCK-01] Isotype IgG1	Application	WB, IP, FC, ICC
human Lck. Lck is a Src-family tyrosine kinase, which is essential for signaling through the T-cell receptor (TCR) complex. Upon TCR triggering, Lck phosphorylates the ITAM motives in its zeta subunits, establishing binding sites for the SH2 domains of the tyro Clonality Monoclonal Concentration 1 mg/ml Clone Designation [LCK-01] Isotype IgG1	Species Reactivity	Human
Product Description through the T-cell receptor (TCR) complex. Upon TCR triggering, Lck phosphorylates the ITAM motives in its zeta subunits, establishing binding sites for the SH2 domains of the tyro Clonality Monoclonal Concentration 1 mg/ml Clone Designation [LCK-01] Isotype IgG1	Immunogen	
Concentration 1 mg/ml Clone Designation [LCK-01] Isotype IgG1	Product Description	through the T-cell receptor (TCR) complex. Upon TCR triggering, Lck phosphorylates the ITAM motives in its zeta subunits, establishing
Clone Designation [LCK-01] Isotype IgG1	Clonality	Monoclonal
Isotype IgG1	Concentration	1 mg/ml
	Clone Designation	[LCK-01]
Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Isotype	lgG1
	Buffer	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Storage 2°C to 8°C	Storage	2°C to 8°C

Antibody Type	Monoclonal Antibody
Application Notes	Flow cytometry: Recommended dilution: 1-5 $\mu g/ml$. Intracellular staining.