

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Anti-Hu CD108 Purified, Clone: [MEM-150], Monoclonal EXB-11-275-C025

Article NameAnti-Hu CD108 Purified, Clone: [MEM-150], MonoclonalBiozol Catalog NumberEXB-11-275-C025Supplier Catalog Number11-275-C025Alternative Catalog NumberEXB-11-275-C025ManufacturerEXBIOCategoryAntikörperApplicationWB, FC, IPSpecies ReactivityHumanImmunogenHPB-ALL human T cell lineProduct DescriptionCD108 (Sema7A) is a GPI-anchored semaphorin family member, which enhances central and peripheral axonal growth and is required for proper axon track formation during ebryogenesis. CD108 also regulates osteoclast differentiation and pre-osteoblastic cClonalityMonoclonalConcentration1 mg/mlClone Designation[MEM-150]IsotypeIgMBufferTris buffered saline (TBS), pH 8.0, 15 mM sodium azideStorage2°C to 8°C		
Supplier Catalog Number 11-275-C025  Alternative Catalog Number EXB-11-275-C025  Manufacturer EXBIO  Category Antikörper  Application WB, FC, IP  Species Reactivity Human  Immunogen HPB-ALL human T cell line  CD108 (Sema7A) is a GPI-anchored semaphorin family member, which enhances central and peripheral axonal growth and is required for proper axon track formation during ebryogenesis. CD108 also regulates osteoclast differentiation and pre-osteoblastic c  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [MEM-150]  Isotype IgM  Buffer Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Article Name	Anti-Hu CD108 Purified, Clone: [MEM-150], Monoclonal
Alternative Catalog Number  EXB-11-275-C025  Manufacturer  EXBIO  Category  Antikörper  Application  WB, FC, IP  Species Reactivity  Human  Immunogen  HPB-ALL human T cell line  CD108 (Sema7A) is a GPI-anchored semaphorin family member, which enhances central and peripheral axonal growth and is required for proper axon track formation during ebryogenesis. CD108 also regulates osteoclast differentiation and pre-osteoblastic c  Clonality  Monoclonal  Concentration  1 mg/ml  Clone Designation  [MEM-150]  Isotype  IgM  Buffer  Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Biozol Catalog Number	EXB-11-275-C025
Manufacturer  EXBIO  Category  Antikörper  Application  WB, FC, IP  Species Reactivity  Human  Immunogen  HPB-ALL human T cell line  CD108 (Sema7A) is a GPI-anchored semaphorin family member, which enhances central and peripheral axonal growth and is required for proper axon track formation during ebryogenesis. CD108 also regulates osteoclast differentiation and pre-osteoblastic c  Clonality  Monoclonal  Concentration  I mg/ml  Clone Designation  [MEM-150]  Isotype  IgM  Buffer  Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Supplier Catalog Number	11-275-C025
Category Antikörper  Application WB, FC, IP  Species Reactivity Human  Immunogen HPB-ALL human T cell line  CD108 (Sema7A) is a GPI-anchored semaphorin family member, which enhances central and peripheral axonal growth and is required for proper axon track formation during ebryogenesis. CD108 also regulates osteoclast differentiation and pre-osteoblastic c  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [MEM-150]  Isotype IgM  Buffer Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Alternative Catalog Number	EXB-11-275-C025
Application  WB, FC, IP  Species Reactivity  Human  HPB-ALL human T cell line  CD108 (Sema7A) is a GPI-anchored semaphorin family member, which enhances central and peripheral axonal growth and is required for proper axon track formation during ebryogenesis. CD108 also regulates osteoclast differentiation and pre-osteoblastic c  Clonality  Monoclonal  Concentration  1 mg/ml  Clone Designation  [MEM-150]  Isotype  IgM  Buffer  Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Manufacturer	EXBIO
Species Reactivity  Human  HPB-ALL human T cell line  CD108 (Sema7A) is a GPI-anchored semaphorin family member, which enhances central and peripheral axonal growth and is required for proper axon track formation during ebryogenesis. CD108 also regulates osteoclast differentiation and pre-osteoblastic c  Clonality  Monoclonal  Concentration  1 mg/ml  Clone Designation  [MEM-150]  Isotype  IgM  Buffer  Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Category	Antikörper
Immunogen       HPB-ALL human T cell line         CD108 (Sema7A) is a GPI-anchored semaphorin family member, which enhances central and peripheral axonal growth and is required for proper axon track formation during ebryogenesis. CD108 also regulates osteoclast differentiation and pre-osteoblastic c         Clonality       Monoclonal         Concentration       1 mg/ml         Clone Designation       [MEM-150]         Isotype       IgM         Buffer       Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Application	WB, FC, IP
Product Description  CD108 (Sema7A) is a GPI-anchored semaphorin family member, which enhances central and peripheral axonal growth and is required for proper axon track formation during ebryogenesis. CD108 also regulates osteoclast differentiation and pre-osteoblastic c  Clonality  Monoclonal  Concentration  1 mg/ml  Clone Designation  [MEM-150]  Isotype  IgM  Buffer  Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Species Reactivity	Human
Product Description which enhances central and peripheral axonal growth and is required for proper axon track formation during ebryogenesis. CD108 also regulates osteoclast differentiation and pre-osteoblastic c  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [MEM-150]  Isotype IgM  Buffer Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Immunogen	HPB-ALL human T cell line
Concentration 1 mg/ml Clone Designation [MEM-150] Isotype IgM Buffer Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Product Description	which enhances central and peripheral axonal growth and is required for proper axon track formation during ebryogenesis. CD108 also
Clone Designation [MEM-150]  Isotype IgM  Buffer Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Clonality	Monoclonal
Isotype IgM  Buffer Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Concentration	1 mg/ml
Buffer Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Clone Designation	[MEM-150]
	Isotype	IgM
Storage 2°C to 8°C	Buffer	Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide
	Storage	2°C to 8°C

Antibody Type	Monoclonal Antibody
Application Notes	Flow cytometry: Recommended dilution: 4 $\mu$ g/ml. Western blotting: Non-reducing conditions.