

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Anti-Hu CD71 Purified, Clone: [MEM-189], Monoclonal EXB-11-353-C100

Article NameAnti-Hu CD71 Purified, Clone: [MEM-189], MonoclonalBiozol Catalog NumberEXB-11-353-C100Supplier Catalog Number11-353-C100Alternative Catalog NumberEXB-11-353-C100ManufacturerEXBIOCategoryAntikörperApplicationFC, IP, WBSpecies ReactivityHumanImmunogenKG1 human acute myelogenous leukaemia cell lineProduct DescriptionCD71 (transferrin receptor) is a type II transmembrane glycoprotein expressed as homodimer in erythroid blood cell line and in activated leukocytes. Upon binding of holotransferrin (complex of transferrin and iron ions), CD71 is internalized by clathClonalityMonoclonalConcentration1 mg/mlClone Designation[MEM-189]IsotypeMouse IgG1BufferTris buffered saline (TBS), pH 8.0, 15 mM sodium azideStorage2°C to 8°C		
Supplier Catalog Number 11-353-C100  Alternative Catalog Number EXB-11-353-C100  Manufacturer EXBIO  Category Antikörper  Application FC, IP, WB  Species Reactivity Human  Immunogen KG1 human acute myelogenous leukaemia cell line  CD71 (transferrin receptor) is a type II transmembrane glycoprotein expressed as homodimer in erythroid blood cell line and in activated leukocytes. Upon binding of holotransferrin (complex of transferrin and iron ions), CD71 is internalized by clath  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [MEM-189]  Isotype Mouse IgG1  Buffer Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Article Name	Anti-Hu CD71 Purified, Clone: [MEM-189], Monoclonal
Alternative Catalog Number EXB-11-353-C100  Manufacturer EXBIO  Category Antikörper  Application FC, IP, WB  Species Reactivity Human  Immunogen KG1 human acute myelogenous leukaemia cell line  CD71 (transferrin receptor) is a type II transmembrane glycoprotein expressed as homodimer in erythroid blood cell line and in activated leukocytes. Upon binding of holotransferrin (complex of transferrin and iron ions), CD71 is internalized by clath  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [MEM-189]  Isotype Mouse IgG1  Buffer Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Biozol Catalog Number	EXB-11-353-C100
Manufacturer  EXBIO  Category  Antikörper  Application  FC, IP, WB  Species Reactivity  Human  KG1 human acute myelogenous leukaemia cell line  CD71 (transferrin receptor) is a type II transmembrane glycoprotein expressed as homodimer in erythroid blood cell line and in activated leukocytes. Upon binding of holotransferrin (complex of transferrin and iron ions), CD71 is internalized by clath  Clonality  Monoclonal  Concentration  I mg/ml  Clone Designation  [MEM-189]  Isotype  Mouse IgG1  Buffer  Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Supplier Catalog Number	11-353-C100
Category Antikörper  Application FC, IP, WB  Species Reactivity Human  Immunogen KG1 human acute myelogenous leukaemia cell line  CD71 (transferrin receptor) is a type II transmembrane glycoprotein expressed as homodimer in erythroid blood cell line and in activated leukocytes. Upon binding of holotransferrin (complex of transferrin and iron ions), CD71 is internalized by clath  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [MEM-189]  Isotype Mouse IgG1  Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Alternative Catalog Number	EXB-11-353-C100
Application FC, IP, WB  Species Reactivity Human  Immunogen KG1 human acute myelogenous leukaemia cell line  CD71 (transferrin receptor) is a type II transmembrane glycoprotein expressed as homodimer in erythroid blood cell line and in activated leukocytes. Upon binding of holotransferrin (complex of transferrin and iron ions), CD71 is internalized by clath  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [MEM-189]  Isotype Mouse IgG1  Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Manufacturer	EXBIO
Species Reactivity  Immunogen  KG1 human acute myelogenous leukaemia cell line  CD71 (transferrin receptor) is a type II transmembrane glycoprotein expressed as homodimer in erythroid blood cell line and in activated leukocytes. Upon binding of holotransferrin (complex of transferrin and iron ions), CD71 is internalized by clath  Clonality  Monoclonal  Concentration  I mg/ml  Clone Designation  [MEM-189]  Isotype  Mouse IgG1  Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Category	Antikörper
Immunogen       KG1 human acute myelogenous leukaemia cell line         Product Description       CD71 (transferrin receptor) is a type II transmembrane glycoprotein expressed as homodimer in erythroid blood cell line and in activated leukocytes. Upon binding of holotransferrin (complex of transferrin and iron ions), CD71 is internalized by clath         Clonality       Monoclonal         Concentration       1 mg/ml         Clone Designation       [MEM-189]         Isotype       Mouse IgG1         Buffer       Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Application	FC, IP, WB
Product Description  CD71 (transferrin receptor) is a type II transmembrane glycoprotein expressed as homodimer in erythroid blood cell line and in activated leukocytes. Upon binding of holotransferrin (complex of transferrin and iron ions), CD71 is internalized by clath  Clonality  Monoclonal  Concentration  1 mg/ml  Clone Designation  [MEM-189]  Isotype  Mouse IgG1  Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Species Reactivity	Human
Product Description  expressed as homodimer in erythroid blood cell line and in activated leukocytes. Upon binding of holotransferrin (complex of transferrin and iron ions), CD71 is internalized by clath  Clonality  Monoclonal  Concentration  1 mg/ml  Clone Designation  [MEM-189]  Isotype  Mouse IgG1  Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Immunogen	KG1 human acute myelogenous leukaemia cell line
Concentration 1 mg/ml Clone Designation [MEM-189] Isotype Mouse IgG1 Buffer Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Product Description	expressed as homodimer in erythroid blood cell line and in activated leukocytes. Upon binding of holotransferrin (complex of transferrin
Clone Designation [MEM-189]  Isotype Mouse IgG1  Buffer Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Clonality	Monoclonal
Isotype Mouse IgG1  Buffer Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Concentration	1 mg/ml
Buffer Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide	Clone Designation	[MEM-189]
The same ca same (125), pit die, 25 mil soaiam aziac	Isotype	Mouse IgG1
Storage 2°C to 8°C	Buffer	Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide
	Storage	2°C to 8°C

Target	CD71
Antibody Type	Monoclonal Antibody
Application Notes	Flow cytometry: Recommended dilution: 2 $\mu$ g/ml. Western blotting: Non-reducing conditions.