

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | Fax: +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-LAT Purified, Clone: [LAT-01], Monoclonal EXB-11-360-C100

Article Name Anti-LAT Purified, Clone: [LAT-01], Monoclonal Biozol Catalog Number EXB-11-360-C100 Supplier Catalog Number 11-360-C100 Alternative Catalog Number EXB-11-360-C100 Manufacturer EXBIO	
Supplier Catalog Number 11-360-C100 Alternative Catalog Number EXB-11-360-C100 Manufacturer EXBIO	
Alternative Catalog Number EXB-11-360-C100 Manufacturer EXBIO	
Manufacturer EXBIO	
Category Antikörper	
Application IP, WB	
Species Reactivity Human	
Immunogen Bacterially produced recombinant polypeptide correspondent entire cytoplasmic domain of human LAT.	onding to the
Product Description LAT (linker for activation of T cells) is a 36-38 kDa doub palmitoylated transmembrane adaptor protein express pre-B cells, NK cells, mast cells and platelets. After immatriggering, LAT becomes multiply tyrosine-phosphoryla.	ed by T cells, nunoreceptor
Clonality Monoclonal	
Concentration 1 mg/ml	
Clone Designation [LAT-01]	
Isotype Mouse IgG1	
Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium	n azide
Storage 2°C to 8°C	

Target	LAT
Antibody Type	Monoclonal Antibody
Application Notes	Immunoprecipitation: Tyrosine-phosphorylated form of LAT is not optimally recognized. Flow cytometry: Recommended dilution: 1-4 $\mu g/ml$, intracellular staining. Tyrosine-phosphorylated form of LAT is not optimally recognized.Western blotting: Recommended dilution: 1-2 $\mu g/ml$.