

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Hu CD38 Purified, Clone: [HIT2], Monoclonal EXB-11-366-C100

Article Name Anti-Hu CD38 Purified, Clone: [HIT2], Monoclonal Biozol Catalog Number EXB-11-366-C100 Supplier Catalog Number 11-366-C100 Alternative Catalog Number EXB-11-366-C100 Manufacturer EXBIO Category Antikörper Application FC, IHC-Fr, IHC-P Species Reactivity Human Immunogen Human thymocytes in foetus CD38 (NAD+ glycohydrolase) is a type II transmembrane glycoprotein able to induce activation, proliferation and differentiation of mature lymphocytes and mediate apoptosis of myeloid and lymphoid progenitor cells. Another role of CD38 is provided by Clonality Monoclonal		
Supplier Catalog Number Alternative Catalog Number EXB-11-366-C100 Manufacturer EXBIO Category Antikörper Application FC, IHC-Fr, IHC-P Species Reactivity Human Immunogen Human thymocytes in foetus CD38 (NAD+ glycohydrolase) is a type II transmembrane glycoprotein able to induce activation, proliferation and differentiation of mature lymphocytes and mediate apoptosis of myeloid and lymphoid progenitor cells. Another role of CD38 is provided by	Article Name	Anti-Hu CD38 Purified, Clone: [HIT2], Monoclonal
Alternative Catalog Number EXB-11-366-C100 Manufacturer EXBIO Category Antikörper Application FC, IHC-Fr, IHC-P Species Reactivity Human Immunogen Human thymocytes in foetus CD38 (NAD+ glycohydrolase) is a type II transmembrane glycoprotein able to induce activation, proliferation and differentiation of mature lymphocytes and mediate apoptosis of myeloid and lymphoid progenitor cells. Another role of CD38 is provided by	Biozol Catalog Number	EXB-11-366-C100
Manufacturer EXBIO Category Antikörper Application FC, IHC-Fr, IHC-P Species Reactivity Human Immunogen Human thymocytes in foetus CD38 (NAD+ glycohydrolase) is a type II transmembrane glycoprotein able to induce activation, proliferation and differentiation of mature lymphocytes and mediate apoptosis of myeloid and lymphoid progenitor cells. Another role of CD38 is provided by	Supplier Catalog Number	11-366-C100
Category Antikörper FC, IHC-Fr, IHC-P Species Reactivity Human Human thymocytes in foetus CD38 (NAD+ glycohydrolase) is a type II transmembrane glycoprotein able to induce activation, proliferation and differentiation of mature lymphocytes and mediate apoptosis of myeloid and lymphoid progenitor cells. Another role of CD38 is provided by	Alternative Catalog Number	EXB-11-366-C100
Application FC, IHC-Fr, IHC-P Species Reactivity Human Human thymocytes in foetus CD38 (NAD+ glycohydrolase) is a type II transmembrane glycoprotein able to induce activation, proliferation and differentiation of mature lymphocytes and mediate apoptosis of myeloid and lymphoid progenitor cells. Another role of CD38 is provided by	Manufacturer	EXBIO
Species Reactivity Human Human thymocytes in foetus CD38 (NAD+ glycohydrolase) is a type II transmembrane glycoprotein able to induce activation, proliferation and differentiation of mature lymphocytes and mediate apoptosis of myeloid and lymphoid progenitor cells. Another role of CD38 is provided by	Category	Antikörper
Immunogen Human thymocytes in foetus CD38 (NAD+ glycohydrolase) is a type II transmembrane glycoprotein able to induce activation, proliferation and differentiation of mature lymphocytes and mediate apoptosis of myeloid and lymphoid progenitor cells. Another role of CD38 is provided by	Application	FC, IHC-Fr, IHC-P
CD38 (NAD+ glycohydrolase) is a type II transmembrane glycoprotein able to induce activation, proliferation and differentiation of mature lymphocytes and mediate apoptosis of myeloid and lymphoid progenitor cells. Another role of CD38 is provided by	Species Reactivity	Human
Product Description glycoprotein able to induce activation, proliferation and differentiation of mature lymphocytes and mediate apoptosis of myeloid and lymphoid progenitor cells. Another role of CD38 is provided by	Immunogen	Human thymocytes in foetus
Clonality Monoclonal	Product Description	glycoprotein able to induce activation, proliferation and differentiation of mature lymphocytes and mediate apoptosis of myeloid and lymphoid progenitor cells. Another role of CD38 is
	Clonality	Monoclonal
Concentration 1 mg/ml	Concentration	1 mg/ml
Clone Designation [HIT2]	Clone Designation	[HIT2]
Isotype Mouse IgG1	Isotype	Mouse IgG1
Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Buffer	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Storage 2°C to 8°C	Storage	2°C to 8°C

Target	CD38
Antibody Type	Monoclonal Antibody
Application Notes	Immunohistochemistry (paraffin sections): Recommended dilution: 10 μg/ml.Flow cytometry: Recommended dilution: 1-4 μg/ml