

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-CSK Purified, Clone: [CSK-04], Monoclonal EXB-11-378-R100

| Anti-CSK Purified, Clone: [CSK-04], Monoclonal |
|--|
| EXB-11-378-R100 |
| 11-378-R100 |
| EXB-11-378-R100 |
| EXBIO |
| Antikörper |
| ICC, IP, WB |
| Human, Mouse |
| Bacterially expressed recombinant fragment of human Csk corresponding to aa 330-450. |
| C-terminal Src kinase (Csk) is a non-receptor protein tyrosine kinase which resembles Src-family kinases, but unlike them it lacks the conserved autophosphorylation site, the regulatory C-terminal tyrosine as well as myristylation and palmitylation |
| Monoclonal |
| [CSK-04] |
| Mouse IgG1 |
| Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide |
| 2°C to 8°C |
| CSK |
| |

| Antibody Type | Monoclonal Antibody |
|-------------------|--|
| Application Notes | Immunocytochemistry: Positive control: HeLa human cervix carcinoma cell line. Western blotting: The recommended dilution of the antibody CSK-04 is 1:250-1:300, weak non-specific band of cca 60 kDa is detected in human cell lines in lower antibody dilutions. Immunoprecipitation: The immunosorbent (CSK-04 antibody covalently bound to beads) immunoprecipitates Csk from cell lysates. |