

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Anti-LIME1 Purified, Clone: [LIME-10], Monoclonal EXB-11-401-C100

|                            |   |
|----------------------------|---|
| Article Name               | Anti-LIME1 Purified, Clone: [LIME-10], Monoclonal   |
| Biozol Catalog Number      | EXB-11-401-C100   |
| Supplier Catalog Number    | 11-401-C100   |
| Alternative Catalog Number | EXB-11-401-C100   |
| Manufacturer               | EXBIO   |
| Category                   | Antikörper  |
| Application                | FC, IHC-P, WB   |
| Species Reactivity         | Human   |
| Immunogen                  | COOH-terminal peptide comprising residues 281-296 of the human LIME conjugated to keyhole limpet hemocyanin.  |
| Product Description        | LIME (Lck-interacting molecule) is a 30 kDa double-palmitoylated protein with unusually basic cytoplasmic domain, expressed by T cells. After ligation of CD4 or CD8 T cell coreceptors, LIME is phosphorylated by Src-family kinases and associates with ... |
| Clonality                  | Monoclonal  |
| Concentration              | 1 mg/ml   |
| Clone Designation          | [LIME-10]   |
| Isotype                    | Mouse IgG2a   |
| Buffer                     | Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide   |
| Storage                    | 2°C to 8°C  |

|                   |  |
|-------------------|--|
| Target            | LIME1  |
| Antibody Type     | Monoclonal Antibody  |
| Application Notes | Immunohistochemistry (paraffin sections): Recommended dilution: 10 µg/ml.Western blotting: Reducing conditions, recommended dilution: 1 µg/ml.Flow cytometry: Recommended dilution: 1-5 µg/ml. |