

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Anti-Hu CD122 APC, Clone: [TU27], Monoclonal EXB-1A-195-T100

|                            |   |
|----------------------------|---|
| Article Name               | Anti-Hu CD122 APC, Clone: [TU27], Monoclonal  |
| Biozol Catalog Number      | EXB-1A-195-T100   |
| Supplier Catalog Number    | 1A-195-T100   |
| Alternative Catalog Number | EXB-1A-195-T100   |
| Manufacturer               | EXBIO   |
| Category                   | Antikörper  |
| Application                | FC  |
| Species Reactivity         | Human   |
| Immunogen                  | TL-Mor cell line  |
| Conjugation                | APC   |
| Product Description        | CD122 (IL-2/IL-15R beta) constitutes together with CD132 (common gamma chain) and with CD25 (IL-2/IL15R alpha) the intermediate (CD122+CD132) and the high affinity (CD122+CD132+CD25) IL-2 and IL-15 receptor complex. CD122 is expressed on NK cells and... |
| Clonality                  | Monoclonal  |
| Clone Designation          | [TU27]  |
| Isotype                    | Mouse IgG1 kappa  |
| Buffer                     | Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide   |
| Storage                    | 2°C to 8°C  |

|                    |   |
|--------------------|---|
| Target             | CD122   |
| Antibody Type      | Monoclonal Antibody   |
| Application Dilute | Flow cytometry: The reagent is designed for analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10 <sup>6</sup> cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests. |
| Application Notes  | Flow cytometry: The reagent is designed for analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10 <sup>6</sup> cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests. |