

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Hu CD11a APC, Clone: [MEM-25], Monoclonal EXB-1A-210-T100

| | |
|----------------------------|---|
| Article Name | Anti-Hu CD11a APC, Clone: [MEM-25], Monoclonal |
| Biozol Catalog Number | EXB-1A-210-T100 |
| Supplier Catalog Number | 1A-210-T100 |
| Alternative Catalog Number | EXB-1A-210-T100 |
| Manufacturer | EXBIO |
| Category | Antikörper |
| Application | FC |
| Species Reactivity | Human |
| Immunogen | Leukocytes from a patient suffering from a LGL-type leukaemia. |
| Conjugation | APC |
| Product Description | CD11a (LFA-1 alpha) together with CD18 constitute leukocyte function-associated antigen 1 (LFA-1), the alphaLbeta2 integrin. CD11a is implicated in activation of LFA-1 complex. LFA-1 is expressed on the plasma membrane of leukocytes in a low-affinity... |
| Clonality | Monoclonal |
| Clone Designation | [MEM-25] |
| Isotype | Mouse IgG1 |
| Buffer | Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide |
| Storage | 2°C to 8°C |

| | |
|--------------------|---|
| Target | CD11a |
| Antibody Type | Monoclonal Antibody |
| Application Dilute | Flow cytometry: The reagent is designed for analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests. |
| Application Notes | Flow cytometry: The reagent is designed for analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests. |