

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Hu CD3 FITC, Clone: [MEM-57], Monoclonal EXB-1F-202-T100

| | |
|----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Article Name | Anti-Hu CD3 FITC, Clone: [MEM-57], Monoclonal |
| Biozol Catalog Number | EXB-1F-202-T100 |
| Supplier Catalog Number | 1F-202-T100 |
| Alternative Catalog Number | EXB-1F-202-T100 |
| Manufacturer | EXBIO |
| Category | Antikörper |
| Application | FC |
| Species Reactivity | Human |
| Immunogen | Human thymocytes and T lymphocytes. |
| Conjugation | FITC |
| Product Description | CD3 complex is crucial in transducing antigen-recognition signals into the cytoplasm of T cells and in regulating the cell surface expression of the TCR complex. T cell activation through the antigen receptor (TCR) involves the cytoplasmic tails of t... |
| Clonality | Monoclonal |
| Clone Designation | [MEM-57] |
| Isotype | Mouse IgG2a kappa |
| Buffer | Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide |
| Storage | 2°C to 8°C |

| | |
|--------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Target | CD3 |
| Antibody Type | Monoclonal Antibody |
| Application Dilute | Flow cytometry: The reagent is designed for analysis of human blood cells using 20 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (2 ml) is sufficient for 100 tests. |
| Application Notes | Flow cytometry: The reagent is designed for analysis of human blood cells using 20 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (2 ml) is sufficient for 100 tests. |