

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Hu CD64 FITC, Clone: [10.1], Monoclonal EXB-1F-644-T100

| | |
|----------------------------|---|
| Article Name | Anti-Hu CD64 FITC, Clone: [10.1], Monoclonal |
| Biozol Catalog Number | EXB-1F-644-T100 |
| Supplier Catalog Number | 1F-644-T100 |
| Alternative Catalog Number | EXB-1F-644-T100 |
| Manufacturer | EXBIO |
| Category | Antikörper |
| Application | FC |
| Species Reactivity | Human, Primate |
| Immunogen | Rheumatoid synovial fluid cells and fibronectin purified human monocytes |
| Conjugation | FITC |
| Product Description | CD64 (FcγRI) is a cell surface receptor for Fc region of IgG. It is composed of specific ligand binding alpha subunit and promiscuous gamma subunit, which is indispensable for tyrosine-based signaling. However, even the alpha subunit can transduc... |
| Clonality | Monoclonal |
| Clone Designation | [10.1] |
| Isotype | Mouse IgG1 kappa |
| Buffer | Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide |

| | |
|--------------------|--|
| Storage | 2°C to 8°C |
| Target | CD64 |
| Antibody Type | Monoclonal Antibody |
| Application Dilute | Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. |
| Application Notes | Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. |