



Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-HLA-G PE, Clone: [87G], Monoclonal

EXB-1P-437-C100

Anti-HLA-G PE, Clone: [87G], Monoclonal
EXB-1P-437-C100
1P-437-C100
EXB-1P-437-C100
EXBIO
Antikörper
FC
Human
HLA-B27 transgenic mice were imunized with H-2 identical murine cells transfected with and expressing genes encoding HLA-G and human beta2-microglobulin.
PE
Human leukocyte antigen G (HLA-G), belonging to MHC class I glycoproteins, plays important roles in both physiological and pathological immunotolerance. It gives an inhibitory signal to cytotoxic T cells, NK cells, monocytes, and some other immune ce
Monoclonal
0.1 mg/ml
[87G]
Mouse IgG2a

Buffer	Stabilizing Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide
Storage	2°C to 8°C
Target	HLA-G
Antibody Type	Monoclonal Antibody
Application Notes	Flow cytometry: Extracellular and intracellular staining, recommended dilution: 2-3 µg/ml, positive control: JEG-3 human choriocarcinoma epithelial cell line.