

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Hu CD193 Alexa Fluor 647, Clone: [5E8], AF647, Monoclonal EXB-A6-161-T100

| | |
|----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Article Name | Anti-Hu CD193 Alexa Fluor 647, Clone: [5E8], AF647, Monoclonal |
| Biozol Catalog Number | EXB-A6-161-T100 |
| Supplier Catalog Number | A6-161-T100 |
| Alternative Catalog Number | EXB-A6-161-T100 |
| Manufacturer | EXBIO |
| Category | Antikörper |
| Application | FC |
| Species Reactivity | Human |
| Immunogen | human CD193 transfectants |
| Conjugation | AF647 |
| Product Description | CD193 / CCR3 is a G-protein coupled receptor for several chemokines, namely CCL11 (eotaxin), CCL26 (eotaxin-3), CCL7 (MCP-4), or CCL5 (RANTES). It is highly expressed on eosinophils and basophils, and is also detected in TH1 and TH2 cells, as well as... |
| Clonality | Monoclonal |
| Clone Designation | [5E8] |
| Isotype | Mouse IgG2b kappa |
| Buffer | Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide |
| Storage | 2°C to 8°C |

| | |
|--------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Target | CD193 |
| Antibody Type | Monoclonal Antibody |
| Application Dilute | Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. |
| Application Notes | Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. |