

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Mouse IgG2a Isotype Control Pacific Blue, Clone: [MOPC-173], Monoclonal EXB-PB-724-C100

|                            |   |
|----------------------------|---|
| Article Name               | Mouse IgG2a Isotype Control Pacific Blue, Clone: [MOPC-173], Monoclonal   |
| Biozol Catalog Number      | EXB-PB-724-C100   |
| Supplier Catalog Number    | PB-724-C100   |
| Alternative Catalog Number | EXB-PB-724-C100   |
| Manufacturer               | EXBIO   |
| Category                   | Antikörper  |
| Application                | FC  |
| Immunogen                  | The transplantable plasmacytoma MOPC-173 was induced by intraperitoneal injection of mineral oils into BALB/c mice. |
| Conjugation                | Pacific Blue  |
| Clonality                  | Monoclonal  |
| Concentration              | 0.1 mg/ml   |
| Clone Designation          | [MOPC-173]  |
| Isotype                    | Mouse IgG2a kappa   |
| Buffer                     | Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide   |
| Storage                    | 2°C to 8°C  |
| Antibody Type              | Monoclonal Antibody   |

|                    |   |
|--------------------|---|
| Application Dilute | Negative control: The reagent is intended as an isotype control to establish the amount of non-specific antibody binding. For your particular experiment, use the same concentration of this control antibody as the recommended working concentration of the   |
| Application Notes  | Negative control: The reagent is intended as an isotype control to establish the amount of non-specific antibody binding. For your particular experiment, use the same concentration of this control antibody as the recommended working concentration of the antigen-specific antibody. Also, when working with prediluted antibodies, dilute the isotype control to the same concentration as is the concentration of the antigen-specific antibody in the prediluted antibody solution you are using. If under particular experimental conditions the background signal of the isotype control is too high (usually when working concentrations of used antibodies are above 10 µg/ml of incubation mixture), change the conditions of your experiment to reduce the background. |