

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Rabbit Antiserum anti-Turkey IgG (H+L)-unconj., MinX none DNA-SEC-182470

|                            |   |
|----------------------------|---|
| Article Name               | Rabbit Antiserum anti-Turkey IgG (H+L)-unconj., MinX none   |
| Biozol Catalog Number      | DNA-SEC-182470  |
| Supplier Catalog Number    | SEC-182470  |
| Alternative Catalog Number | DNA-SEC-182470  |
| Manufacturer               | dianova   |
| Host                       | Rabbit  |
| Category                   | Antikörper  |
| Application                | ELISA,IHC,WB  |
| Species Reactivity         | Turkey  |
| Immunogen                  | Turkey IgG whole molecule   |
| Conjugation                | Unconjugated  |
| Format                     | Antiserum   |
| Target Specificity         | IgG (H+L)   |
| Cross-Adsorption (MinX)    | no cross-adsorbtion   |
| Product Description        | Anti-Turkey IgG (H&L) Antibody generated in rabbit detects reactivity to Turkey IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bac... |
| Clonality                  | Polyclonal  |

|                    |   |
|--------------------|---|
| Concentration      | 90 mg/mL  |
| Isotype            | Ig  |
| Buffer             | 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2   |
| Purity             | This product was prepared from monospecific antiserum by a delipidation and defibrination. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-rabbit serum, Turkey IgG and Turkey Serum. |
| Form               | Lyophilized   |
| Formula            | 10 mM NaPO <sub>4</sub> , 150 mM NaCl, pH 7.2, lyophilisate, 0.01% NaN <sub>3</sub>   |
| Target             | Turkey  |
| Antibody Type      | Secondary Antibody  |
| Application Dilute | ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000   |
| Application Notes  | Anti-Turkey IgG (H&L) Antibody are ideal for ELISA, western blotting, Immunohistochemistry, as well as other antibody detection methods.  |