

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit Anti-Swine IgG F(ab)2 Antibody Peroxidase Conjugated - 214-4304, HRP, Polyclonal DNA-SEC-182633

Article Name	Rabbit Anti-Swine IgG F(ab)2 Antibody Peroxidase Conjugated - 214-4304, HRP, Polyclonal
Biozol Catalog Number	DNA-SEC-182633
Supplier Catalog Number	DNA-SEC-182633
Alternative Catalog Number	DNA-SEC-182633
Manufacturer	dianova
Host	Rabbit
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Porcine
Immunogen	Swine IgG F(ab)2 fragment
Conjugation	HRP
Format	IgG
Target Specificity	IgG (F(ab')2)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	Anti-Swine IgG F(ab)2 Antibody generated in rabbit recognizes the dimeric Fab portion of the swine IgG molecule. Swine IgG F(ab)2 is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme pepsin under controlle...

Clonality	Polyclonal
Concentration	10.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum, Swine IgG, Swine IgG F(ab)2 and Swine Serum. No reaction was observed against Swine IgG F(c).
Form	Lyophilized
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin
Target	Swine
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:10,000 - 1:50,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, Western Blot Dilution: 1:1,000 - 1:10,000
Application Notes	This product has been assayed against 1.0 ug of Swine IgG in a standard capture ELISA using ABTS (2,2-azino-bis-[3-ethylbenzthiazoline-6-sulfonic acid]) code ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:27,000 to 1:110,000 of the reconstitution concentration is suggested for this product.