

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit F(ab)2 Anti-Bovine IgG F(ab)2 Antibody Fluorescein Conjugated - 301-4204, FITC, Polyclonal DNA-SEC-182648

Article Name	Rabbit F(ab)2 Anti-Bovine IgG F(ab)2 Antibody Fluorescein Conjugated - 301-4204, FITC, Polyclonal
Biozol Catalog Number	DNA-SEC-182648
Supplier Catalog Number	DNA-SEC-182648
Alternative Catalog Number	DNA-SEC-182648
Manufacturer	dianova
Host	Rabbit
Category	Antikörper
Application	FLISA,FACS,IF
Species Reactivity	Bovine
Immunogen	Bovine IgG F(ab)2 fragment
Conjugation	FITC
Format	F(ab')2
Target Specificity	IgG (F(ab')2)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	F(ab)2 Anti-Bovine IgG F(ab)2 Antibody generated in rabbit detects Bovine F(ab)2. Representing approximately 75% of serum immunoglobulins, IgG is the most abundant antibody isotype found in the circulation. IgG molecules are synthesized a...

Clonality	Polyclonal
Concentration	10.0 mg/mL
Isotype	Ig
Buffer	0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product is a F(ab)2 fragment of an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation, ion exchange chromatography and pepsin digestion followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Rabbit Serum, Bovine IgG, Bovine IgG F(ab)2 and Bovine Serum. No reaction was observed against Bovine IgG F(c), anti-Rabbit IgG F(c) or anti-Pepsin.
Form	Lyophilized
Formula	10 mM NaPO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3
Target	Bovine
Antibody Type	Secondary Antibody
Application Dilute	FLISA Dilution: 1:10,000 - 1:50,000, Flow Cytometry Dilution: 1:500 - 1:2,500, Fluorochrome Protein Value: 3.1, IF Microscopy Dilution: 1:1,000 - 1:5,000
Application Notes	Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.