

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Rabbit F(ab)2 anti-Chicken IgG (H+L)-FITC, MinX none DNA-SEC-182657

Article Name	Rabbit F(ab)2 anti-Chicken IgG (H+L)-FITC, MinX none
Biozol Catalog Number	DNA-SEC-182657
Supplier Catalog Number	SEC-182657
Alternative Catalog Number	DNA-SEC-182657
Manufacturer	dianova
Host	Rabbit
Category	Antikörper
Application	FLISA,FACS,IF
Species Reactivity	Gallus
Immunogen	Chicken IgG whole molecule
Conjugation	FITC
Format	F(ab')2
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial pla...
Clonality	Polyclonal

Concentration	10.0 mg/mL
Isotype	Ig
Buffer	0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product is a F(ab') <sub>2</sub> fragment of an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation, ion exchange chromatography and pepsin digestion followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-fluorescein, anti-Rabbit Serum, Chicken IgG and Chicken Serum. No reaction was observed against anti-Rabbit IgG F(c) or anti-Pepsin.
Form	Lyophilized
Formula	10 mM NaPO <sub>4</sub> , 150 mM NaCl, pH 7.2, lyophilisate, 0.01% Thimerosal
Target	Chicken
Antibody Type	Secondary Antibody
Application Dilute	FLISA Dilution: 1:10,000 - 1:50,000, Flow Cytometry Dilution: 1:500 - 1:2,500, Fluorochrome Protein Value: 3.2, IF Microscopy Dilution: 1:1,000 - 1:5,000
Application Notes	Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.