

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Rabbit F(ab)2 anti-Goat IgG (F(ab)2)-unconj., MinX none DNA-SEC-182667

Article Name	Rabbit F(ab)2 anti-Goat IgG (F(ab)2)-unconj., MinX none
Biozol Catalog Number	DNA-SEC-182667
Supplier Catalog Number	SEC-182667
Alternative Catalog Number	DNA-SEC-182667
Manufacturer	dianova
Host	Rabbit
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Goat
Immunogen	Goat IgG F(ab)2 fragment
Conjugation	Unconjugated
Format	F(ab')2
Target Specificity	IgG (F(ab')2)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	F(ab)2 Anti-Goat IgG F(ab)2 Antibody generated in rabbit detects Goat F(ab)2. Representing approximately 75% of serum immunoglobulins, IgG is the most abundant antibody isotype found in the circulation. IgG molecules are synthesized and secreted by p...
Clonality	Polyclonal

Concentration	10.0 mg/mL
Isotype	Ig
Buffer	0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product is a F(ab') <sub>2</sub> fragment of IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation, ion exchange chromatography and pepsin digestion followed by chromatographic separation and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Goat IgG, Goat IgG F(ab') <sub>2</sub> and Goat Serum. No reaction was observed against Goat IgG F(c), anti-Rabbit IgG F(c) or anti-Pepsin.
Form	Lyophilized
Formula	10 mM NaPO <sub>4</sub> , 150 mM NaCl, pH 7.2, lyophilisate, 0.01% NaN <sub>3</sub>
Target	Goat
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000
Application Notes	Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maximum amount of reagent required to stain 1 x 10 <sup>6</sup> cells in flow cytometry is approximately 1.0 µg of antibody. Lesser amounts of reagent may be sufficient for staining. Optimal titers for other applications should be determined by the researcher. As a general guideline dilutions of 1:100 to 1:250 should be suitable for most applications.