

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Rabbit F(ab)2 anti-Goat IgG (H+L)-FITC, MinX none DNA-SEC-182668

Article Name	Rabbit F(ab)2 anti-Goat IgG (H+L)-FITC, MinX none
Biozol Catalog Number	DNA-SEC-182668
Supplier Catalog Number	SEC-182668
Alternative Catalog Number	DNA-SEC-182668
Manufacturer	dianova
Host	Rabbit
Category	Antikörper
Application	FLISA,FACS,IF
Species Reactivity	Goat
Immunogen	Goat IgG whole molecule
Conjugation	FITC
Format	F(ab')2
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	F(ab)2 Anti-Goat IgG Fluorescein Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for use in certain immunoche...
Clonality	Polyclonal

Concentration	10.0 mg/mL
Isotype	Ig
Buffer	0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product is a F(ab') <sub>2</sub> fragment of an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation, ion exchange chromatography and pepsin digestion followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Rabbit Serum, Goat IgG and Goat Serum. No reaction was observed against anti-Rabbit IgG F(c) or anti-Pepsin.
Form	Lyophilized
Formula	10 mM NaPO <sub>4</sub> , 150 mM NaCl, pH 7.2, lyophilisate, 0.01% NaN <sub>3</sub>
Target	Goat
Antibody Type	Secondary Antibody
Application Dilute	FLISA Dilution: 1:10,000 - 1:50,000, Flow Cytometry Dilution: 1:500-1:2,500, Fluorochrome Protein Value: 4.0, IF Microscopy Dilution: 1:1,000-1:5,000
Application Notes	Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.