

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit F(ab)2 anti-Horse IgG (H+L)-unconj., MinX none DNA-SEC-182683

| | |
|----------------------------|---|
| Article Name | Rabbit F(ab)2 anti-Horse IgG (H+L)-unconj., MinX none |
| Biozol Catalog Number | DNA-SEC-182683 |
| Supplier Catalog Number | SEC-182683 |
| Alternative Catalog Number | DNA-SEC-182683 |
| Manufacturer | dianova |
| Host | Rabbit |
| Category | Antikörper |
| Application | ELISA,IHC,WB |
| Species Reactivity | Equine |
| Immunogen | Horse IgG whole molecule |
| Conjugation | Unconjugated |
| Format | F(ab')2 |
| Target Specificity | IgG (H+L) |
| Cross-Adsorption (MinX) | no cross-adsorbtion |
| Product Description | F(ab)2 Anti-Horse IgG Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for use in certain immunochemical techn... |
| Clonality | Polyclonal |

| | |
|--------------------|---|
| Isotype | Ig |
| Buffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Purity | This product is a F(ab') ₂ fragment of IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation, ion exchange chromatography and pepsin digestion followed by chromatographic separation and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Horse IgG and Horse Serum. No reaction was observed against anti-Rabbit IgG F(c) or anti-Pepsin. |
| Form | Lyophilized |
| Formula | 20 mM K ₃ PO ₄ , 150 mM NaCl, pH 7.2, lyophilisate, Azide/BSA free |
| Target | Horse |
| Antibody Type | Secondary Antibody |
| Application Dilute | ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000-1:5,000, Western Blot Dilution: 1:2,000-1:10,000 |
| Application Notes | Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maximum amount of reagent required to stain 1 x 10 ⁶ cells in flow cytometry is approximately 1.0 µg of antibody. Lesser amounts of reagent may be sufficient for staining. Optimal titers for other applications should be determined by the researcher. As a general guideline dilutions of 1:100 to 1:250 should be suitable for most applications. |