

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat F(ab)2 anti-Human IgG (H+L)-HRPO, MinX none DNA-SEC-182689

Article Name	Goat F(ab)2 anti-Human IgG (H+L)-HRPO, MinX none
Biozol Catalog Number	DNA-SEC-182689
Supplier Catalog Number	SEC-182689
Alternative Catalog Number	DNA-SEC-182689
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Human
Immunogen	Human IgG whole molecule
Conjugation	HRPO
Format	F(ab')2
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	F(ab)2 Anti-Human IgG (H&L) Peroxidase Antibody generated in goat detects immunoglobulin g from human, both heavy and light chains of the antibody molecule are present. Each IgG has two antigen binding sites. Representing approximately 75% of serum i...
Clonality	Polyclonal

Concentration	10.0 mg/mL
Isotype	Ig
Buffer	0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product is a F(ab') ₂ fragment of an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation, ion exchange chromatography and pepsin digestion followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Human IgG and Human Serum. No reaction was observed against anti-Goat IgG F(c) or anti-Pepsin.
Form	Lyophilized
Formula	10 mM NaPO ₄ , 150 mM NaCl, pH 7.2, lyophilisate, 0.01% Gentamicin
Target	Human
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:10,000 - 1:50,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, Western Blot Dilution: 1:1,000 - 1:10,000
Application Notes	F(ab) ₂ Anti-Human IgG (H&L) Peroxidase secondary antibody reagents are ideal for ELISA, western blotting, Immunohistochemistry, Fluorescence Microscopy, Flow Cytometry as well as other antibody detection methods.