

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Rabbit IgG anti-Bovine IgG (H+L)-unconj., MinX none DNA-SEC-182728

| Auticle Nieure             | DILL'IL COULD IN LOCKHALLY   |
|----------------------------|--|
| Article Name               | Rabbit IgG anti-Bovine IgG (H+L)-unconj., MinX none  |
| Biozol Catalog Number      | DNA-SEC-182728   |
| Supplier Catalog Number    | SEC-182728   |
| Alternative Catalog Number | DNA-SEC-182728   |
| Manufacturer               | dianova  |
| Host                       | Rabbit   |
| Category                   | Antikörper   |
| Application                | ELISA,IHC,WB   |
| Species Reactivity         | Bovine   |
| Immunogen                  | Bovine IgG whole molecule  |
| Conjugation                | Unconjugated   |
| Format                     | IgG  |
| Target Specificity         | IgG (H+L)  |
| Cross-Adsorption (MinX)    | no cross-adsorbtion  |
| Product Description        | Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglu |
| Clonality                  | Polyclonal   |

| Concentration      | 2.0 mg/mL  |
|--------------------|--|
| Isotype            | Ig   |
| Buffer             | 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2  |
| Purity             | Anti-Bovine IgG (H&L) Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Bovine IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Bovine IgG and Bovine Serum. |
| Form               | Liquid (sterile filtered)  |
| Formula            | 10 mM NaPO4,150 mM NaCl,pH 7,2,sterile filtered,0,01% NaN3   |
| Target             | Bovine   |
| Antibody Type      | Secondary Antibody   |
| Application Dilute | ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000  |
| Application Notes  | Anti-Bovine IgG (H&L) Antibody has been tested by ELISA and is suitable for immunoblotting (western or dot blot), ELISA, immunoelectron microscopy and immunohistochemistry as well as other antibody based enzymatic assays requiring lot-to-lot consistency.   |