

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit IgG anti-Bovine IgG (H+L)-HRPO, MinX none DNA-SEC-182734

Article Name	Rabbit IgG anti-Bovine IgG (H+L)-HRPO, MinX none
Biozol Catalog Number	DNA-SEC-182734
Supplier Catalog Number	SEC-182734
Alternative Catalog Number	DNA-SEC-182734
Manufacturer	dianova
Host	Rabbit
Category	Antikörper
Application	ELISA,IHC
Species Reactivity	Bovine
Immunogen	Bovine IgG whole molecule
Conjugation	HRPO
Format	IgG
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglu
Clonality	Polyclonal

Concentration	2.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	Anti-Bovine IgG antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Bovine IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum, Bovine IgG and Bovine Serum.
Form	Lyophilized
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin
Target	Bovine
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:10,000 - 1:50,000, Immunohistochemistry Dilution: 1:500 - 1:2,500
Application Notes	Anti Bovine IgG antibody has been assayed against $1.0~\mu g$ of Bovine IgG in a standard capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) as a substrate for 30 minutes at room temperature. A working dilution of $1:1,000$ to $1:5,000$ of the reconstitution concentration is suggested for this product. Specific conditions for reactivity should be optimized by the end user.