

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit IgG anti-Chicken IgG (F(ab)2)-Alk. Phos., MinX none DNA-SEC-182801

Article Name	Rabbit IgG anti-Chicken IgG (F(ab)2)-Alk. Phos., MinX none
Biozol Catalog Number	DNA-SEC-182801
Supplier Catalog Number	SEC-182801
Alternative Catalog Number	DNA-SEC-182801
Manufacturer	dianova
Host	Rabbit
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Gallus
Immunogen	Chicken IgG F(ab)2 fragment
Conjugation	Alk. Phos.
Format	IgG
Target Specificity	IgG (F(ab')2)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	Anti-Chicken IgG F(ab)2 Antibody generated in rabbit is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme pepsin under controlled conditions of temperature, time and pH. F(ab)2 molecules lack the Fc portio...
Clonality	Polyclonal

Concentration	1.0 mg/mL
Isotype	Ig
Buffer	0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Chicken IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Alkaline Phosphatase (calf intestine), anti-Rabbit Serum, Chicken IgG, Chicken IgG F(ab')2 and Chicken Serum. No reaction was observed against Chicken IgG F(c).
Form	Liquid (sterile filtered)
Formula	50 mM TrisHCl,150 mM NaCl,1 mM MgCl,0,1 mM ZnCl,50% (v/v) Glycerol,pH 8,0,sterile filtered,0,01% NaN3
Target	Chicken
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:3,500, Immunohistochemistry Dilution: 1:200 - 1:1,000, Western Blot Dilution: 1:500 - 1:2,500
Application Notes	Anti-Chicken IgG F(ab')2 antibody is suitable for immunoblotting (western or dot blot), ELISA, immunoelectron microscopy and immunohistochemistry as well as other antibody based enzymatic assays requiring lot-to-lot consistency.