

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | Fax: +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit IgG anti-Goat IgG (H+L)-FITC, MinX none DNA-SEC-182846

Article Name	Rabbit IgG anti-Goat IgG (H+L)-FITC, MinX none
Biozol Catalog Number	DNA-SEC-182846
Supplier Catalog Number	SEC-182846
Alternative Catalog Number	DNA-SEC-182846
Manufacturer	dianova
Host	Rabbit
Category	Antikörper
Application	FLISA,FACS,IF
Species Reactivity	Goat
Immunogen	Goat IgG whole molecule
Conjugation	FITC
Format	IgG
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	Anti-Goat IgG Fluorescein Antibody generated in rabbit detects goat IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as we
Clonality	Polyclonal

Concentration	2.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	Anti-Goat Secondary Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Rabbit Serum, Goat IgG and Goat Serum.
Form	Lyophilized
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3
Target	Goat
Antibody Type	Secondary Antibody
Application Dilute	FLISA Dilution: 1:10,000 - 1:50,000, Flow Cytometry Dilution: 1:500 - 1:2,500, Fluorochrome Protein Value: 3.73, IF Microscopy Dilution: 1:1,000 - 1:5,000
Application Notes	Secondary antibody reagents are ideal for western blotting, Immunohistochemistry, ELISA, Fluorescence Microscopy, Flow Cytometry as well as other antibody detection methods.