

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit IgG anti-Goat IgG (F(ab)2)-Alk. Phos., MinX Hu, ALP, Polyclonal , AP DNA-SEC-182863

Article Name	Rabbit IgG anti-Goat IgG (F(ab)2)-Alk. Phos., MinX Hu, ALP, Polyclonal , AP
Biozol Catalog Number	DNA-SEC-182863
Supplier Catalog Number	SEC-182863
Alternative Catalog Number	DNA-SEC-182863
Manufacturer	dianova
Host	Rabbit
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Goat
Immunogen	Goat IgG F(ab)2 fragment
Conjugation	Alk. Phos.
Format	IgG
Target Specificity	IgG (F(ab')2)
Cross-Adsorption (MinX)	Human
Product Description	Anti-Goat IgG F(ab)2 Alkaline Phosphatase Antibody generated in rabbit is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme pepsin under controlled conditions of temperature, time and pH. F(ab)2 molecules ...

Clonality	Polyclonal
Concentration	1.0 mg/mL
Isotype	Ig
Buffer	0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Alkaline Phosphatase (calf intestine), anti-Rabbit Serum, Goat IgG, Goat IgG F(ab') ₂ and Goat Serum. No reaction was observed against Goat IgG F(c) or Human Serum Proteins.
Form	Liquid (sterile filtered)
Formula	50 mM TrisHCl, 150 mM NaCl, 1 mM MgCl, 0.1 mM ZnCl, 50% (v/v) Glycerol, pH 8.0, sterile filtered, 0.01% NaN ₃
Target	Goat
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:2,000 - 1:10,000, Immunohistochemistry Dilution: 1:200 - 1:1,000, Western Blot Dilution: 1:500 - 1:2,500
Application Notes	This product has been assayed against 1.0 µg of Goat IgG in a standard capture ELISA using pNPP p-nitrophenyl phosphate code # NPP-10 as a substrate for 30 minutes at room temperature. A working dilution of 1:2,000 to 1:8,000 is suggested for this product.