

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Donkey IgG anti-Goat IgG (H+L)-Biotin, MinX Hm,Rb,Rt,Ck,Gp,Ho,Ms, Polyclonal DNA-SEC-182890

Article Name	Donkey IgG anti-Goat IgG (H+L)-Biotin, MinX Hm,Rb,Rt,Ck,Gp,Ho,Ms, Polyclonal
Biozol Catalog Number	DNA-SEC-182890
Supplier Catalog Number	SEC-182890
Alternative Catalog Number	DNA-SEC-182890
Manufacturer	dianova
Host	Donkey
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Goat
Immunogen	Anti-Goat IgG (H&L) was produced by repeated immunization with goat IgG whole molecule in donkey.
Conjugation	Biotin
Format	IgG
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	Hamster (all),Rabbit,Rat,Gallus,Guinea pig,Equine,Mouse
Product Description	Anti-Goat IgG Biotin Antibody generated in donkey detects goat IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as...

Clonality	Polyclonal
Concentration	1.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	Anti-Goat IgG (H&L) Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Donkey Serum, Goat IgG and Goat Serum. No reaction was observed against Chicken, Guinea Pig, Horse, Hamster, Mouse, Rabbit and Rat Serum Proteins.
Form	Lyophilized
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3
Target	Goat
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:600,000, Fluorochrome Protein Value: 10-20, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:60,000
Application Notes	Goat IgG (H&L) Antibody is suitable for immunoblotting, ELISA, immunohistochemistry, immunomicroscopy as well as other antibody based assays using streptavidin or avidin conjugates requiring lot-to-lot consistency.