

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Chicken IgG anti-Goat IgG (H+L)-Alk. Phos., MinX none DNA-SEC-182895

| | |
|----------------------------|---|
| Article Name | Chicken IgG anti-Goat IgG (H+L)-Alk. Phos., MinX none |
| Biozol Catalog Number | DNA-SEC-182895 |
| Supplier Catalog Number | SEC-182895 |
| Alternative Catalog Number | DNA-SEC-182895 |
| Manufacturer | dianova |
| Host | Gallus |
| Category | Antikörper |
| Application | ELISA,IHC,WB |
| Species Reactivity | Goat |
| Immunogen | Goat IgG whole molecule |
| Conjugation | Alk. Phos. |
| Format | IgG |
| Target Specificity | IgG (H+L) |
| Cross-Adsorption (MinX) | no cross-adsorbtion |
| Product Description | Anti-Goat IgG Alkaline Phosphatase Antibody generated in chicken detects goat IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacte... |
| Clonality | Polyclonal |

| | |
|--------------------|---|
| Concentration | 1.0 mg/mL |
| Isotype | Ig |
| Buffer | 0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0 |
| Purity | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Alkaline Phosphatase (calf intestine), anti-Chicken Serum, Goat IgG and Goat Serum. |
| Form | Liquid (sterile filtered) |
| Formula | 50 mM TrisHCl,150 mM NaCl,1 mM MgCl ₂ ,0,1 mM ZnCl ₂ ,50% (v/v) Glycerol,pH 8,0,sterile filtered,0,01% NaN ₃ |
| Target | Goat |
| Antibody Type | Secondary Antibody |
| Application Dilute | ELISA Dilution: 1:2,000 - 1:10,000, Immunohistochemistry Dilution: 1:200 - 1:1,000, Western Blot Dilution: 1:500 - 1:2,500 |
| Application Notes | This product has been assayed against 1.0 ug of Goat IgG in a standard capture ELISA using pNPP (p-nitrophenyl phosphate) code # NPP-10 as a substrate for 30 minutes at room temperature. A working dilution of 1:500 to 1:2,000 of the stated concentration is suggested for this product. |